PARAMETERIZED WASSERSTEIN GRADIENT FLOW

YIJIE JIN *, SHU LIU[†], HAO WU[‡], XIAOJING YE[§], AND HAOMIN ZHOU[¶]

Abstract. We develop a fast and scalable numerical approach to solve Wasserstein gradient flows (WGFs), which is particularly suitable for high-dimensional cases. Our approach is to use general reduced-order models, like deep neural networks, to parameterize the push-forward maps such that they can push a simple reference density to the one solving the given WGF. The new dynamical system is called parameterized WGF (PWGF), and it is defined on the finite-dimensional parameter space equipped with a pullback Wasserstein metric. Our numerical scheme can approximate the solutions of WGFs for general energy functionals effectively, without requiring spatial discretization or nonconvex optimization procedures, thus avoiding some limitations of classical numerical methods and more recent deep-learning based approaches. A comprehensive analysis of the approximation errors measured by Wasserstein distance is also provided in this work. Numerical experiments show promising computational efficiency and verified accuracy on a variety of WGF examples using our approach.

Key words. Wasserstein gradient flow; Fokker-Planck equation; Porous medium equation; Deep neural networks; Numerical analysis.

1. Introduction. Wasserstein gradient flow (WGF) is a powerful tool for understanding and analyzing density evolution processes. In the seminal work [20] by Jordan, Kinderlehrer and Otto, they showed that Fokker-Planck equation (FPE) is essentially the gradient flow of the relative entropy functional under the Wasserstein metric. Since then, WGFs have shown extensive applications in optimal transport theory, optimization problems, Fokker-Planck equation, porous medium equation, and more [11, 29, 32]. However, numerical computation of general WGFs remains a challenging problem, especially when the state space is of high dimension. Furthermore, it is often desirable to find a sampler that generates samples following the solution of WGF rather than the actual density function solving WGF in many real-world statistics and machine learning applications.

In this paper, we focus on the numerical computation of WGF to address the aforementioned issues. Let \mathcal{M} be a smooth manifold without boundary. For simplicity, we assume $\mathcal{M} = \mathbb{R}^d$ throughout the present work, while generalization to general manifolds is straightforward. We also omit subscript \mathcal{M} for all integrals unless otherwise noted. Denote the density function space defined on \mathcal{M} as

(1.1)
$$\mathcal{P}(\mathcal{M}) = \Big\{ \rho : \mathcal{M} \to \mathbb{R} \colon \int \rho(x) dx = 1, \ \rho(x) \ge 0, \ \int |x|^2 \rho(x) \ dx < \infty \Big\}.$$

Suppose $\mathcal{P}(\mathcal{M})$ is equipped with the Wasserstein-2 distance [23, 32]:

(1.2)
$$W(\rho_1, \rho_2) = \left(\inf_{\pi \in \Pi(\rho_1, \rho_2)} \iint |x - y|^2 d\pi(x, y)\right)^{1/2}$$

for any $\rho_1, \rho_2 \in \mathcal{P}(\mathcal{M})$, where $\Pi(\rho_1, \rho_2)$ is the joint density space with ρ_1 and ρ_2 as marginals. Then $\mathcal{P}(\mathcal{M})$ becomes an infinite-dimensional Riemannian manifold with W inducing its Riemannian metric, and a WGF can be written in the following general form:

(1.3)
$$\frac{\partial \rho}{\partial t} = -\operatorname{grad}_W \mathcal{F}(\rho), \quad \rho(0, x) = \rho_0(x),$$

where $x \in \mathcal{M}$, ρ_0 is a given initial probability density, $\mathcal{F} : \mathcal{P}(\mathcal{M}) \to \mathbb{R}$ is some energy functional defined on $\mathcal{P}(\mathcal{M})$, and grad_W stands for the gradient of functionals on $\mathcal{P}(\mathcal{M})$ with respect to the Wasserstein metric. Many well-known probability evolution equations, such as the Fokker-Planck equation and porous medium equation, are essentially WGF (1.3) with specific choices of \mathcal{F} .

In recent years, there has been a surge of interest to compute WGF (1.3) numerically. In [20], a proximal point algorithm is used to solve the Fokker-Planck equation as a special case of WGF. Later, various numerical methods have been developed to solve the WGF; see [9, 14, 30, 31, 33]. Among them, some use classical methods such as finite difference [25] and finite element method [6]. They are applicable to solving WGF with low dimension d. This limitation is due to their spatial discretization and hence they suffer the issue known

^{*}School of Mathematics, Georgia Institute of Technology, Atlanta, GA, USA (yijiejin@gatech.edu).

[†]Department of Mathematics, University of California, Los Angles, CA, USA(shuliu@math.ucla.edu).

[‡]School of Mathematics, Georgia Institute of Technology, Atlanta, GA, USA (hwu406@gmail.com).

[§]Department of Mathematics and Statistics, Georgia State University, Atlanta, GA, USA(xye@gsu.edu).

[¶]School of Mathematics, Georgia Institute of Technology, Atlanta, GA, USA(hmzhou@gatech.edu).

as "curse-of-dimensionality" as the number of unknowns increases exponentially fast in terms of problem dimension d. Sampling-based approaches [14, 30] can be used to solve WGF in high dimension, however, the computation for the minimization and energy functional evaluation used in those algorithms may become time-consuming.

The goal of this work is to develop a fast computational scheme of (1.3) in high-dimensional settings. Similar to [14, 29, 30], we seek for a time-dependent push-forward map to generate samples whose density function solves (1.3). However, compared to these existing methods, our approach features several notable differences: (i) We use a *new pullback Wasserstein metric* that can be computed efficiently while still maintaining the accuracy up to the same order as the results obtained in [29]; (ii) Our algorithm can be used to compute general WGF with a large class of (possibly nonlinear) \mathcal{F} ; (iii) We parameterize the push-forward map using neural ordinary differential equation [10] and show that it is computationally convenient to handle functional of probability densities in our case. As a result, we obtain an approximate solution to the WGF as well as a sample generator of the solution distribution; and (iv) Our numerical scheme does not need any spatial discretization and thus is scalable for problems defined on high-dimensional spaces (e.g., $d \ge 15$). Moreover, our method does not need to solve any nonconvex optimization problem as in typical deep neural network training. Instead, it only requires fast matrix-vector products where the vector size is determined by the number of parameters in the selected push-forward map.

2. Related work. The Fokker-Planck equation is first interpreted as a special case of WGF in [20]. It is shown that the Fokker-Planck equation can be viewed as the gradient flow on the Wasserstein manifold of a special energy functional, which consists of a linear potential energy and a relative entropy. For general (possibly nonlinear) energy functionals, one can apply similar derivations to obtain their corresponding WGFs. As WGFs are essentially time-evolution partial differential equations (PDEs), classical numerical methods such as finite difference and finite element methods can be applied with proper modifications [3, 8, 9, 27, 33]. However, the application of these classical methods is limited to low-dimensional cases due to the curse of dimensionality.

In recent years, sampling-based approaches have been proposed as a promising alternative solution method [14, 19, 24, 29, 30]. Many of these methods start from the numerical scheme used in [20] known as the Jordan-Kinderlehrer-Otto (JKO) scheme, which is a proximal algorithm in the optimization context. By utilizing the Benamou-Brenier formula, we know that for any time step, the optimal push-forward map in the JKO scheme can be expressed as the gradient of some convex function. Motivated by this fact, several recent works are devoted to solving the WGF by approximating the optimal push-forward map by neural networks T_{θ} . However, there are two main challenges in this approach: one is the evaluation of potential energy \mathcal{F} , especially when \mathcal{F} involves the density function ρ explicitly. As mentioned in [26], this problem can be addressed by introducing an extra optimization procedure. However, such a computation is generally expensive; the other is the computational challenge for solving the JKO scheme, which is an optimization problem on the density manifold.

In contrast to those existing methods, our method is mostly motivated by [29], in which the authors develop the parameterization to solve FPE. This approach exploits WGFs in the parameter space, in which the parameterize functions, such as deep neural networks, are used to parameterize the push-forward maps. We design a numerical scheme for the parameter dynamics. The main challenge in our formulation is the computation of the Wasserstein gradient on the parameter manifold. We show that it can be effectively approximated once introducing the new pullback Wasserstein metric as discussed in (3.17). Furthermore, we employ several powerful neural networks, such as the normalizing flow [21] and continuous normalizing flow [10], which are both convenient and efficient in evaluating the energy functionals involving push-forward densities and their gradients.

In addition to the aforementioned literatures, it is worth mentioning that the study on the Wasserstein information matrix as the metric tensor defined on the parameter manifold has been introduced in [28]; Some numerical analysis results on 1-dimensional neural projected WGFs have been reported in [41]; Studies on WGFs on the manifold of Gaussian distributions have been conducted in [11, 40] and the references therein.

Furthermore, a series of deep-learning methods [1, 7, 13, 16] have been composed to compute general timeevolution PDEs. Compared with these references which directly approximate the PDE solutions via neural networks, our algorithm is developed on the space of push-forward maps and utilizes the geometric principle of the WGFs. Our treatment respects the essential properties of WGFs such as positivity, conservation of mass, and energy dissipation: the first two properties are naturally preserved on push-forward models; the energy dissipation property is justified via our numerical experiments demonstrated in Section 4. **3.** Parameterization of Wasserstein gradient flow. In this section, we briefly review the derivation of WGF and several examples of WGF including Fokker-Planck and porous media equations, as well as the pullback Wasserstein metric in parameter space. Then we present our approach to solve WGF numerically and provide a comprehensive analysis of the proposed scheme.

3.1. Background on Wasserstein gradient flow. We denote the tangent space of $\mathcal{P}(\mathcal{M})$ at ρ by

$$\mathcal{T}_{\rho}\mathcal{P}(\mathcal{M}) = \Big\{ \sigma \in C^{\infty}(\mathcal{M}) \colon \int \sigma(x) \, dx = 0 \Big\}.$$

For any specific $\rho \in \mathcal{P}(\mathcal{M})$, the Wasserstein metric tensor g^W is a positive definite bilinear form on the tangent bundle $\mathcal{TP}(\mathcal{M}) = \{(\rho, \sigma) : \rho \in \mathcal{P}(\mathcal{M}), \sigma \in \mathcal{T}_{\rho}\mathcal{P}(\mathcal{M})\}$ defined by:

$$g^{W}(\rho)(\sigma_{1},\sigma_{2}) = \int \nabla \Phi_{1}(x) \cdot \nabla \Phi_{2}(x)\rho(x) \, dx, \quad \forall \sigma_{i} \in \mathcal{T}_{\rho}\mathcal{P}(M), \ i = 1,2$$

where $\sigma_i = -\nabla \cdot (\rho \nabla \Phi_i)$ for i = 1, 2. Suppose $\mathcal{F} \colon \mathcal{P}(\mathcal{M}) \to \mathbb{R}$ is a smooth functional defined on $\mathcal{P}(\mathcal{M})$, its Riemannian gradient over $(\mathcal{P}(\mathcal{M}), g^W)$ is given as follows:

(3.1)
$$\operatorname{grad}_{W}\mathcal{F}(\rho) = g^{W}(\rho)^{-1} \left(\frac{\delta\mathcal{F}}{\delta\rho}\right)(x) = -\nabla \cdot \left(\rho(x)\nabla \frac{\delta\mathcal{F}}{\delta\rho}(x)\right)$$

where $\frac{\delta \mathcal{F}}{\delta \rho}$ is the first variation of \mathcal{F} in the L^2 sense. The Wasserstein gradient flow (WGF) of \mathcal{F} is given by

(3.2)
$$\frac{\partial \rho}{\partial t} = -\text{grad}_W \mathcal{F}(\rho).$$

Many well-known equations can be formulated as WGF with selected energy functionals. Here are three examples.

EXAMPLE 3.1 (Fokker-Planck equation). Let $V \in C^2(M)$ be a given potential function and $\rho_*(x) = \frac{1}{Z_D}e^{-V(x)/D}$ be the corresponding Gibbs distribution, where D > 0 is a fixed constant and $Z_D = \int e^{-\frac{V(x)}{D}} dx$ is the normalization constant. Suppose $\mathcal{F}(\rho)$ is the relative entropy with respect to ρ_* scaled by D, i.e.,

(3.3)
$$\mathcal{F}(\rho) = D \mathcal{H}_{KL}(\rho \| \rho_*) \quad \text{where} \quad \mathcal{H}_{KL}(\rho \| \rho_*) := \left(\int \frac{1}{D} V(x) \rho(x) + \rho(x) \log \rho(x) \, dx \right) + \log Z_D.$$

Then the WGF (3.2) with \mathcal{F} defined in (3.3) becomes the Fokker-Planck equation:

(3.4)
$$\frac{\partial \rho}{\partial t} = -\operatorname{grad}_W \mathcal{F}(\rho) = \nabla \cdot (\rho \nabla V) + D\nabla \cdot (\rho \nabla \log \rho) = \nabla \cdot (\rho \nabla V) + D\Delta \rho.$$

The relative entropy $\mathcal{H}_{KL}(\rho \| \rho_*)$ is closely related to the Fisher information defined as

(3.5)
$$\mathcal{I}(\rho|\rho_*) = \int \left\|\nabla \log\left(\frac{\rho(x)}{\rho_*(x)}\right)\right\|^2 \rho(x) \, dx.$$

It provides an upper bound to \mathcal{H}_{KL} and further guarantees the uniform convergence of the dynamics (3.4) as shown in the following theorem, which will be used in our error estimation later.

THEOREM 3.2 (Holley–Stroock perturbation [18]). Suppose the potential V can be decomposed as $V = U + \phi$ where $\nabla^2 U \succeq KI$ for some K > 0 and $\phi \in L^{\infty}$. Denote $osc(\phi) := \sup \phi - \inf \phi < \infty$. Then the following logarithmic Sobolev inequality holds

(3.6)
$$\mathcal{H}_{KL}(\rho \| \rho_*) \leqslant \frac{e^{osc(\phi)}}{K} \mathcal{I}(\rho | \rho_*)$$

for any $\rho \in \mathcal{P}(M)$. Assume ρ solves equation (3.4) with initial value $\rho(0, \cdot) = \rho_0(x)$, then

(3.7)
$$\mathcal{H}_{KL}(\rho_t \| \rho_*) \leqslant \mathcal{H}_{KL}(\rho_0 \| \rho_*) e^{-\frac{DK}{osc(\phi)}t}, \quad \forall t > 0.$$

EXAMPLE 3.3 (Porous medium equation). Consider the energy functional \mathcal{F} defined by

(3.8)
$$\mathcal{F}(\rho) = \frac{1}{m-1} \int \rho^m(x) \, dx.$$

for an integer m > 1. The corresponding WGF becomes the porous medium equation (PME), which is a nonlinear heat equation given by

(3.9)
$$\frac{\partial \rho}{\partial t} = \Delta(\rho^m).$$

There are many applications of PME in physical problems, for example, heat transfer or diffusion, gas flow [36], and flow of reactive fluids in porous media [22].

EXAMPLE 3.4 (Aggregation model). Consider the interaction energy functional

(3.10)
$$\mathcal{F}(\rho) = \frac{1}{2} \iint J(|x-y|)\rho(x)\rho(y) \, dx \, dy$$

which is used to model the interactive behavior of swarm of particles [15]. Here J is an interaction kernel consisting of repulsive and attractive parts. A typical choice for J is

(3.11)
$$J(x) = \frac{|x|^a}{a} - \frac{|x|^b}{b}, \quad x \in \mathcal{M}.$$

where a > b > 0 are positive constants. Its corresponding WGF is

(3.12)
$$\frac{\partial \rho}{\partial t} = \nabla \cdot (\rho \nabla (J * \rho))$$

We note that Keller-Siegel system is a special case of aggregation model [5]. We remark that the WGF (3.2) can be associated with a particle-level dynamic, as given in the following proposition.

PROPOSITION 3.5. [29] Assume $X \in \mathcal{M}$ is a random process and solves the following equation,

(3.13)
$$\dot{\boldsymbol{X}} = -\nabla_{\boldsymbol{X}} \frac{\delta}{\delta \rho} \mathcal{F}(\rho(\boldsymbol{X}), \boldsymbol{X})$$

where ρ is the density of **X**. Then ρ solves the WGF (3.2).

The particle-level dynamic (3.13) provides a physical interpretation of WGF. More importantly, we will use it to establish the error bound of our approximation using parameterized WGF below.

3.2. Push-forward maps and parameterized Fokker-Planck equation. In [29], the authors developed a method to parameterize push-forward maps in order to approximately solve the FPE (3.4).

Fix any reference probability distribution λ that is absolutely continuous with respect to the standard Lebesgue measure μ on \mathcal{M} , we use $\rho = d\lambda/d\mu$, the Radon-Nikodym derivative of λ with respect to μ , to denote the reference density determined by λ . Then a push-forward map $T : \mathbb{R}^d \to \mathbb{R}^d$ induces a new probability density $T_{\sharp}\rho$ on \mathbb{R}^d :

$$\int_E T_{\sharp} \varrho(x) \, d\mu(x) = \int_{T^{-1}(E)} \varrho(z) \, d\mu(z) = \int_{T^{-1}(E)} d\lambda(z) = \lambda(T^{-1}(E)) \quad \text{ for all measurable } E \subset \mathbb{R}^d,$$

where $T^{-1}(E)$ is the pre-image of E. Hereafter we use dx instead of $d\mu(x)$ to reduce notation complexity.

Let us take T as parameterized map, namely for any $\theta \in \Theta$, $T_{\theta} : \mathbb{R}^d \to \mathbb{R}^d$ is a parametric function with parameter θ . Here Θ , as a subset of \mathbb{R}^n , is called the *parameter space*, where n is the number of parameters of T_{θ} (i.e., the dimension of θ). Typical examples of T_{θ} include Fourier expansion, finite element approximation, and (deep) neural networks.

The map $T_{\sharp}: \Theta \to \mathcal{P}$ given by $\theta \mapsto T_{\theta \sharp} \varrho$ naturally defines an immersion from Θ to the probability manifold \mathcal{P} . Collecting all parameterized distributions together, i.e.,

$$\mathcal{P}_{\Theta} = \left\{ \rho_{\theta} = T_{\theta \sharp} \varrho : \ \theta \in \Theta \right\},$$

we obtain a finite-dimensional submanifold \mathcal{P}_{Θ} of \mathcal{P} . We can define the tangent space of \mathcal{P}_{Θ} at each $\theta = (\theta_1, \ldots, \theta_n) \in \Theta \subset \mathbb{R}^n$ as $T_{\rho_{\theta}} \mathcal{P}_{\Theta} = \operatorname{span}\{\frac{\partial \rho_{\theta}}{\partial \theta_1}, \cdots, \frac{\partial \rho_{\theta}}{\partial \theta_n}\}$. The tangent bundle is then $\mathcal{T}\mathcal{P}_{\Theta} = \bigcup_{\theta \in \Theta}\{\rho_{\theta}\} \times T_{\rho_{\theta}}\mathcal{P}_{\Theta}$. On the other hand, the cotangent space $T^*_{\rho_{\theta}}\mathcal{P}_{\Theta}$ is the dual space of $T_{\rho_{\theta}}\mathcal{P}_{\Theta}$, and the cotangent bundle is $\mathcal{T}^*\mathcal{P}_{\Theta} = \bigcup_{\theta \in \Theta}\{\rho_{\theta}\} \times T^*_{\rho_{\theta}}\mathcal{P}_{\Theta}$.

Note that \mathcal{P}_{Θ} is a finite-dimensional subset of \mathcal{P} , hence we can pull back the Wasserstein metric g^W to \mathcal{P}_{Θ} . It is shown in [29] that the pullback Wasserstein metric $G(\theta) = T^*_{\theta \sharp} g^W$ on \mathcal{P} is given by

(3.14)
$$G(\theta) = \int \nabla \Psi \circ T_{\theta}(z) \nabla \Psi \circ T_{\theta}(z)^{\top} d\lambda(z).$$

where $\Psi_{\theta} = (\psi_1, \cdots, \psi_n)^{\top}$, and ψ_j is solved from the following elliptic equation,

(3.15)
$$\nabla \cdot (\rho_{\theta} \nabla \psi_j(x)) = \nabla \cdot \left(\rho_{\theta} \frac{\partial T_{\theta}}{\partial \theta_j} \circ T_{\theta}^{-1}(x)\right).$$

The parameterized version of FPE on parameter manifold Θ can be expressed as

(3.16)
$$\dot{\theta} = -G(\theta)^{-1} \nabla_{\theta} F(\theta),$$

where $F(\theta) := \mathcal{F}(\rho_{\theta})$ and \mathcal{F} is the scaled relative entropy defined in (3.3).

If directly computing the parameterized FPE (3.16), the cost is intractable in higher dimensions because it requires solving *n* elliptic equations (3.15) to obtain $G(\theta)$. To alleviate this difficulty, authors in [29] derived a minimax formulation that can advance the dynamics without solving the elliptic equations. Although the cost becomes manageable, it can still be high if the spatial dimension *d* is not small. In this work, we advocate a new, yet naturally derived pullback metric \hat{G} to replace *G*, and develop a computational framework to solve general WGF. Details shall be provided in the next subsection.

3.3. Parameterized WGF with a new pullback Wasserstein metric. Following the study in [39], we replace $G(\theta)$ by a new pullback Wasserstein metric defined as

(3.17)
$$\widehat{G}(\theta) = \int \partial_{\theta} T_{\theta}(z)^{\top} \partial_{\theta} T_{\theta}(z) \ d\lambda(z)$$

For any energy functional \mathcal{F} with smooth variation $\frac{\delta \mathcal{F}}{\delta \rho}$, the corresponding parameterized WGF is given by

(3.18)
$$\dot{\theta} = -\widehat{G}(\theta)^{\dagger} \nabla_{\theta} F(\theta),$$

where $\widehat{G}(\theta)^{\dagger}$ is the Penrose-Moore pseudo inverse of $\widehat{G}(\theta)$.

The motivation of introducing $\widehat{G}(\theta)$ is three-folded:

- The definition is directly inspired by (3.14) and (3.15). In the 1-dimensional case, the new metric \hat{G} coincides with the exact matrix G, and the proof is shown in [29].
- A deeper motivation is influenced by [32]. The geometry of $(\Theta, \widehat{G}(\theta))$ is an isometric embedding into the flat Riemannian space $(\mathcal{O}, g^{\mathcal{O}})$ introduced in [32]. Specifically, let $\varrho \in P(\mathcal{M})$ be a fixed density function and \mathcal{O} the set of diffeomorphisms $T : \mathcal{M} \to \mathcal{M}$, and define the Riemannian metric $g^{\mathcal{O}}$ on \mathcal{O} at T by $g^{\mathcal{O}}(T)(v_1, v_2) := \int v_1 v_2 \rho \, dx$ for any $v_1, v_2 \in \mathcal{T}_T \mathcal{O}$. Here $\mathcal{T}_T \mathcal{O}$ is the tangent space of \mathcal{O} at T, which is set to be the space of all vector fields on \mathcal{M} for every T in [32]. Then it is shown that the mapping $\Pi : \theta \mapsto \rho_{\theta}$ yields an isometric submersion from $(\mathcal{O}, g^{\mathcal{O}})$ to $(P(\mathcal{M}), g)$, where g is the 2-Wasserstein metric. Consider a (deep) neural network structure T_{θ} with parameter θ such that $T_{\theta} : \mathcal{M} \to \mathcal{M}$ is a diffeomorphism for every $\theta \in \Theta$ and $T_{\cdot}(\cdot) \in C^1(\Theta \times \mathcal{M}; \mathbb{R})$. Consider the map $\mathscr{T} : \Theta \to \mathcal{O}$ defined by $\theta \mapsto \mathscr{T}(\theta) := T_{\theta}$, we define the pullback metric on Θ as

(3.19)
$$\widehat{G}(\theta) = \mathscr{T}^* g^{\mathcal{O}}(T_\theta)$$

where \mathscr{T}^* is the pullback operation induced by \mathscr{T} . To obtain $\widehat{G}(\theta)$, consider any curve $\{\theta(t)\}_{-\epsilon \leq t \leq \epsilon}$ for some $\epsilon > 0$ passing through $\theta(0)$ at t = 0 and denote $\dot{\theta}(0) = \frac{d}{dt}\theta(t)|_{t=0}$. By the definitions of the

pullback operation and the metric $g^{\mathcal{O}}$ above, we have

$$\begin{aligned} \widehat{G}(\theta(0))(\dot{\theta}(0), \dot{\theta}(0)) &= g^{\mathcal{O}}(T_{\theta}) \left(\frac{d}{dt} T_{\theta(t)}|_{t=0}, \frac{d}{dt} T_{\theta(t)}|_{t=0} \right) \\ &= \int \left(\dot{\theta}(0)^{\top} \partial_{\theta} T_{\theta(0)}(z)^{\top} \partial_{\theta} T_{\theta(0)}(z) \dot{\theta}(0) \right) \varrho(z) \, dz \\ &= \dot{\theta}(0)^{\top} \left(\int \partial_{\theta} T_{\theta(0)}(z)^{\top} \partial_{\theta} T_{\theta(0)}(z) \, d\lambda(z) \right) \dot{\theta}(0), \end{aligned}$$

which implies (3.17) since $\theta(0)$ and $\dot{\theta}(0)$ are arbitrary. The matrix $\hat{G}(\theta) = (\hat{G}(\theta)_{ij})_{1 \leq i,j \leq m}$ has components

(3.20)
$$\widehat{G}(\theta)_{ij} = \sum_{k=1}^{d} \int \partial_{\theta_i} T_{\theta}^{(k)}(z) \cdot \partial_{\theta_j} T_{\theta}^{(k)}(z) \, d\lambda(z)$$

where $T_{\theta}^{(k)} : \mathcal{M} \to \mathbb{R}$ is the k-th component of $T_{\theta} : \mathcal{M} \to \mathcal{M}$. This is also discussed in [38].

• Compared with (3.14), the new metric is more computationally efficient. Directly evaluating the metric tensor G requires solving n different elliptic PDEs where n is the number of the parameters in the push-forward map T_{θ} , and m can be large if we choose T_{θ} to be a deep neural network. To circumvent the computation complexity challenge, a bi-level minimization scheme was proposed in [29]. However, it may still be computationally expensive to solve such optimization problems in general. We demonstrate the difference in the computation time in the experiment for computing the solution of the Fokker-Planck equation in Section 5.1.

The gradient $\nabla_{\theta} F(\theta)$ can be evaluated through the following formulation as derived in [39],

(3.21)
$$\nabla_{\theta} F(\theta) = \int \partial_{\theta} T_{\theta}(z)^{\top} \nabla_{X} \frac{\delta}{\delta \rho} \mathcal{F}(T_{\theta \sharp} \varrho(\cdot), \cdot) \circ T_{\theta}(z) \ d\lambda(z).$$

Following [39], the formula (3.21) suggests that $\nabla_{\theta} F(\theta)$ is in the range of $\widehat{G}(\theta)$ even when \widehat{G} is not invertible. Therefore, the right hand side of (3.18) is always well defined. For convenience, we still call (3.18) the parameterized WGF, or again PWGF for short.

It is interesting to consider the particle level dynamics corresponding to (3.18). For a fix $z_0 \in \mathcal{M}$, $\mathbf{Y} = T_{\theta}(z_0)$ is the push-forward point in \mathcal{M} . When θ varies as a function of t according to (3.18), $\mathbf{Y}(t)$ forms a curve in \mathcal{M} , and it satisfies

(3.22)
$$\dot{\boldsymbol{Y}} = \partial_{\theta} T_{\theta}(z_0) \dot{\theta} = -\partial_{\theta} T_{\theta}(z_0) \widehat{G}(\theta)^{\dagger} \nabla_{\theta} F(\theta), \quad \boldsymbol{Y}_0 = T_{\theta(0)}(z_0).$$

To further investigate the properties of parameterized particle dynamics (3.22), we introduce the kernel operator \mathcal{K} : for any $f \in L^2(\mathcal{M}; \mathcal{M}, \lambda)$, $\mathcal{K}_{\theta}[f] \in L^2(\mathcal{M}; \mathcal{M}, \lambda)$ is defined as

(3.23)
$$\mathcal{K}_{\theta}[f](\cdot) = \partial_{\theta} T_{\theta}(\cdot) \widehat{G}(\theta)^{\dagger} \int \partial_{\theta} T_{\theta}(z)^{\top} f(z) \ d\lambda(z) = \int K_{\theta}(\cdot, z) f(z) \ d\lambda(z),$$

where the kernel matrix $K_{\theta}(z', z) \in \mathbb{R}^{d \times d}$ is defined by

(3.24)
$$K_{\theta}(z',z) := \partial_{\theta} T_{\theta}(z') \widehat{G}(\theta)^{\dagger} \partial_{\theta} T_{\theta}(z)^{\top}.$$

It is shown that $\mathcal{K}_{\theta}[f]$ is the orthogonal projection of f onto the tangent space spanned by $\partial_{\theta}T_{\theta}$ [39]. Combining (3.21), (3.22) and (3.23), we obtain

$$\dot{\boldsymbol{Y}} = -\partial_{\theta} T_{\theta}(z_{0}) \widehat{\boldsymbol{G}}(\theta)^{\dagger} \int \partial_{\theta} T_{\theta}(z)^{\top} \nabla_{X} \frac{\delta}{\delta\rho} \mathcal{F}(T_{\theta\sharp}\varrho(\cdot), \cdot) \circ T_{\theta}(z) \ d\lambda(z)$$

$$= -\mathcal{K}_{\theta} [\nabla_{X} \frac{\delta}{\delta\rho} \mathcal{F}(T_{\theta\sharp}\varrho(\cdot), \cdot) \circ T_{\theta}](z_{0})$$

$$= -\mathcal{K}_{\theta} [\nabla_{X} \frac{\delta}{\delta\rho} \mathcal{F}(T_{\theta\sharp}\varrho(\cdot), \cdot) \circ T_{\theta}](T_{\theta}^{-1}(\boldsymbol{Y})).$$

The particle-level dynamics (3.25) is determined by the time-evolving push-forward map T_{θ} . We will use it to derive an upper bound on the error of PWGF (3.18) in the next section.

3.4. Error bounds in Wasserstein metric. We provide an error analysis of the proposed PWGF formulation. Specifically, we establish an upper bound on the difference, measured by Wasserstein distance, between the approximation $\rho_{\theta(t)}(\cdot) = T_{\theta(t)\sharp}\varrho(\cdot)$ where $\theta(t)$ solves PWGF and the true solution $\rho(t, \cdot)$ to the original WGF. We consider the error analysis in two scenarios: the first variation $\frac{\delta \mathcal{F}}{\delta \rho}$ is Lipschitz continuous and the case of Fokker-Planck equation where $\frac{\delta \mathcal{F}}{\delta \rho}$ is not Lipschitz continuous.

3.4.1. Error analysis with Lipschitz continuity assumption. In this subsection, we assume the following condition on \mathcal{F} holds.

Assumption 1. There exists a constant $C_{\mathcal{F}} > 0$ such that for any two push-forward maps T and \widetilde{T} there is

$$(3.26) \qquad \int \left| \nabla_X \frac{\delta}{\delta\rho} \mathcal{F}(T_{\sharp} \varrho(\cdot), \cdot) \circ T(z) - \nabla_X \frac{\delta}{\delta\rho} \mathcal{F}(\widetilde{T}_{\sharp} \varrho(\cdot), \cdot) \circ \widetilde{T}(z) \right|^2 \, d\lambda(z) \leqslant C_{\mathcal{F}} \int |T(z) - \widetilde{T}(z)|^2 \, d\lambda(z).$$

The accuracy of our approach depends on the representation power of the parameterized push-forward map T_{θ} . We use δ_0 to characterize the representation error as below.

DEFINITION 3.6. Define the projection error as

$$\delta_{0} = \sup_{\theta \in \Theta} \min_{\xi \in \mathcal{T}_{\theta}^{*} \Theta} \left\{ \int \left| \nabla \frac{\delta}{\delta \rho} \mathcal{F}(T_{\theta \sharp} \varrho(\cdot), \cdot) \circ T_{\theta}(z) - \partial_{\theta} T_{\theta}(z) \xi \right|^{2} d\lambda(z) \right\}$$

$$(3.27) \qquad \qquad = \sup_{\theta \in \Theta} \left\{ \int \left| \nabla \frac{\delta}{\delta \rho} \mathcal{F}(T_{\theta \sharp} \varrho(\cdot), \cdot) \circ T_{\theta}(z) - \mathcal{K}_{\theta} [\nabla \frac{\delta}{\delta \rho} \mathcal{F}(T_{\theta \sharp} \varrho(\cdot), \cdot) \circ T_{\theta}](z) \right|^{2} d\lambda(z) \right\}.$$

This error is essentially the difference between $\frac{\delta \mathcal{F}(\rho_{\theta})}{\delta \rho}$ and its orthogonal projection onto the tangent space $T_{\rho_{\theta}}\mathcal{P}(\mathcal{M})$. In the definition, we use the fact that the operator \mathcal{K}_{θ} is the orthogonal projection. Under Assumption 1, we can show that the density ρ_{θ} approximates the true solution with guaranteed Wasserstein-2 error stated in the next theorem.

THEOREM 3.7. Suppose Assumption 1 holds for \mathcal{F} , θ is solved from (3.18) with initial value $\theta(0)$, and ρ is solved from (1.3). Then the Wasserstein-2 distance between the push-forward density $\rho_{\theta(t)}(\cdot)$ obtained by PWGF and the true density $\rho(t, \cdot)$ satisfies

(3.28)
$$W_2^2(\rho_{\theta(t)}(\cdot), \rho(t, \cdot)) \leqslant e^{(1+2C_{\mathcal{F}})t} \epsilon_{\rho(0)} + \frac{2\delta_0}{1+2C_{\mathcal{F}}} \left(e^{(1+2C_{\mathcal{F}})t} - 1 \right).$$

where $\epsilon_{\rho(0)} = W_2^2(\rho_{\theta(0)}, \rho_0)$ is the initial approximation error.

Proof. Assume \boldsymbol{X} is solved from (3.13) and \boldsymbol{Y} is solved from (3.22). Suppose the Monge map from $\rho_{\theta(0)}$ to ρ_0 is given by ω and we assume the random variables $\boldsymbol{X}, \boldsymbol{Y}$ are coupled via $\boldsymbol{X}(0) = \omega(\boldsymbol{Y}(0))$. We continue from the calculation in (3.25),

$$\begin{split} \dot{\boldsymbol{Y}} &= -\mathcal{K}_{\theta} \left[\frac{\delta}{\delta\rho} \mathcal{F}(T_{\theta \sharp} \varrho(\cdot), \cdot) \circ T_{\theta} \right] (z) \\ &= -\frac{\delta}{\delta\rho} \mathcal{F}(T_{\theta \sharp} \varrho(\cdot), \cdot) \circ T_{\theta}(z) + \left(\frac{\delta}{\delta\rho} \mathcal{F}(T_{\theta \sharp} \varrho(\cdot), \cdot) \circ T_{\theta}(z) - \mathcal{K}_{\theta} \left[\frac{\delta}{\delta\rho} \mathcal{F}(T_{\theta \sharp} \varrho(\cdot), \cdot) \circ T_{\theta} \right] (z) \right). \end{split}$$

Denote $E(t) = \mathbb{E} \| \boldsymbol{X} - \boldsymbol{Y} \|^2$, we compute,

$$(3.29) \quad \frac{d}{dt}E(t) = 2\mathbb{E}\left[(\boldsymbol{X} - \boldsymbol{Y}) \cdot (\dot{\boldsymbol{X}} - \dot{\boldsymbol{Y}})\right] \leq 2\sqrt{\mathbb{E}\|\boldsymbol{X} - \boldsymbol{Y}\|^2}\sqrt{\mathbb{E}\|\dot{\boldsymbol{X}} - \dot{\boldsymbol{Y}}\|^2} \leq \mathbb{E}\|\boldsymbol{X} - \boldsymbol{Y}\|^2 + \mathbb{E}\|\dot{\boldsymbol{X}} - \dot{\boldsymbol{Y}}\|^2.$$

Notice that X is a push-forward of X(0) through the dynamics (3.13), we have

$$\mathbb{E}\|\dot{\boldsymbol{X}} - \dot{\boldsymbol{Y}}\|^{2} = \int \left| -\left(\frac{\delta}{\delta\rho}\mathcal{F}(\rho(x), x) - \frac{\delta}{\delta\rho}\mathcal{F}(T_{\theta\sharp}\varrho(\cdot), \cdot) \circ T_{\theta}(z)\right) - \left(\frac{\delta}{\delta\rho}\mathcal{F}(T_{\theta\sharp}\varrho(\cdot), \cdot) \circ T_{\theta}(z) - \mathcal{K}_{\theta}\left[\frac{\delta}{\delta\rho}\mathcal{F}(T_{\theta\sharp}\varrho(\cdot), \cdot) \circ T_{\theta}\right](z)\right) \right|^{2} d\lambda(z)$$

$$\leq 2\delta_{0} + 2C_{\mathcal{F}} \mathbb{E}\|\boldsymbol{X} - \boldsymbol{Y}\|^{2}.$$

(3.30)

Plugging (3.30) into (3.29), we obtain

(3.31)
$$\frac{d}{dt}E(t) \leq 2\delta_0 + (1+2C_{\mathcal{F}})E(t)$$

By Grönwall's inequality, we have

(3.32)
$$E(t) \leqslant e^{(1+2C_{\mathcal{F}})t}E(0) + \frac{2\delta_0}{1+2C_{\mathcal{F}}}\left(e^{(1+2C_{\mathcal{F}})t} - 1\right).$$

On the other hand, we have

(3.33)
$$W_2^2(T_{\theta \sharp} \varrho, \rho) = W_2^2(\text{Law}(\boldsymbol{Y}), \text{Law}(\boldsymbol{X})) \leqslant \mathbb{E} \|\boldsymbol{Y} - \boldsymbol{X}\|^2 = E(t).$$

Combining (3.32) and (3.33) yields (3.28).

3.4.2. Asymptotic Analysis for Parametrized Wasserstein Gradient Flow. For many examples, finding $C_{\mathcal{F}}$ could be a challenging task. However, if there exists a Gibbs solution ρ_* to the WGF satisfying Polyak-Lojasiewicz inequality, then we can show uniform convergence of ρ_{θ} to the target density ρ_* , as stated in the following theorem:

THEOREM 3.8. Assume that ρ_* is the Gibbs distribution, and the functional $\mathcal{F}(\rho)$ satisfies certain Polyak-Lojasiewicz inequality as follows

(3.34)
$$\frac{1}{\zeta} \int_{\mathbb{R}^d} \left| \nabla_X \frac{\delta \mathcal{F}}{\delta \rho}(\rho, x) \right|^2 \rho(x) \ dx \ge \mathcal{F}(\rho) - \mathcal{F}(\rho_*)$$

where ζ is a positive constant. Then

(3.35)
$$\mathcal{F}(\rho_{\theta(t)}) - \mathcal{F}(\rho_*) \leqslant \frac{\delta_0}{\zeta} (1 - e^{-\zeta t}) + \mathcal{F}(\rho_{\theta(0)}) e^{-\zeta t}$$

where δ_0 is defined in (3.27).

Proof. Denote $f(z) = \nabla_X \frac{\delta}{\delta\rho} \mathcal{F}(T_{\theta \sharp} \varrho(\cdot), \cdot) \circ T_{\theta}(z)$. Then by the property of orthogonal projection operator \mathcal{K}_{θ} we have $\int (I - \mathcal{K}_{\theta}) [f](z)^{\top} \mathcal{K}_{\theta}[f](z) \ d\lambda(z) = 0$. Furthermore,

(3.36)
$$\int f(z)^{\top} \mathcal{K}_{\theta}[f](z) \ d\lambda(z) = \int \mathcal{K}_{\theta}[f](z)^{\top} \mathcal{K}_{\theta}[f](z) \ d\lambda(z) + \int (I - \mathcal{K}_{\theta})[f](z)^{\top} \mathcal{K}_{\theta}[f](z) \ d\lambda(z)$$
$$= \int \mathcal{K}_{\theta}[f](z)^{\top} \mathcal{K}_{\theta}[f](z) \ d\lambda(z)$$
$$= \int |f(z)|^{2} \ d\lambda(z) - \int |(I - \mathcal{K}_{\theta})[f](z)|^{2} \ d\lambda(z).$$

where the last equality is due to the Pythagoras theorem. Following the definition of δ_0 in (3.27), we have

(3.37)
$$\int |(I - \mathcal{K}_{\theta})[f](z)|^2 d\lambda(z) \leq \delta_0$$

By inequality (3.34), we have

(3.38)
$$\int |f(z)|^2 d\lambda(z) \ge \zeta \mathcal{F}(\rho)$$

We deduce that

(3.39)

$$\begin{split} \frac{d}{dt}\mathcal{F}(\rho_{\theta}) &= \nabla_{\theta}F(\theta)\cdot\dot{\theta} \\ &= -\nabla_{\theta}F(\theta)^{\top}\widehat{G}(\theta)^{\dagger}\nabla_{\theta}F(\theta) \\ &= -\int \nabla_{X}\frac{\delta}{\delta\rho}\mathcal{F}(T_{\theta\sharp}\varrho(\cdot),\cdot)\circ T_{\theta}(z)^{\top}\partial_{\theta}T_{\theta}(z)\widehat{G}(\theta)^{\dagger}\nabla_{\theta}F(\theta) \ d\lambda(z) \\ &= -\int \nabla_{X}\frac{\delta}{\delta\rho}\mathcal{F}(T_{\theta\sharp}\lambda(\cdot),\cdot)\circ T_{\theta}(z)^{\top}\mathcal{K}_{\theta}[\nabla_{X}\frac{\delta}{\delta\rho}\mathcal{F}(T_{\theta\sharp}\lambda(\cdot),\cdot)\circ T_{\theta}](z) \ d\lambda(z) \\ &= -\int f(z)^{\top}\mathcal{K}_{\theta}[f](z) \ d\lambda(z), \end{split}$$

where the first equality is due to $F(\theta) := \mathcal{F}(\rho_{\theta})$, the second equality is obtained by the formulation of PWGF in (3.18), the third equality uses the substitution of first $\nabla_{\theta} F(\theta)$ using (3.21), the fourth equality is because of the definition of \mathcal{K}_{θ} in (3.23), and the fifth equality is by the definition of f at the beginning of this proof. We also observe that

(3.40)
$$-\int f(z)^{\top} \mathcal{K}_{\theta}[f](z) \ d\lambda(z) \leq \delta_{0} - \int |f(z)|^{2} \ d\lambda(z) \leq \delta_{0} - \zeta \mathcal{F}(\rho_{\theta}),$$

where the first inequality is obtained by combining (3.36) and (3.37), the second equality is due to (3.38). Combining (3.39) and (3.40) yields

(3.41)
$$\frac{d}{dt}F(\theta) \leqslant \delta_0 - \zeta F(\theta).$$

Applying the Grönwall inequality to (3.41) yields the claimed estimate (3.35).

We would like to mention that the bound obtained in Theorem 3.8 is similar to the results reported in [29] except that they use different pullback metrics, namely we use $\hat{G}(\theta)$ while $G(\theta)$ is considered in [29].

4. Numerical method. The PWGF (3.18) can be solved based on standard numerical integrators. We showcase how to implement this using the forward Euler scheme, whereas the idea can be easily generated to other methods such as Runge-Kutta 4th-order method, predictor-corrector method, and those with variable step sizes. In forward Euler scheme, we can disretize the time t as $\{hl: l = 0, 1, ...\}$. Let θ^l be the approximation of $\theta(hl)$ of PWGF (3.18), then we can compute θ^{l+1} given θ^{l} by solving

(4.1)
$$\frac{\theta^{l+1} - \theta^l}{h} = -\widehat{G}(\theta^l)^{\dagger} \nabla_{\theta} F(\theta^l),$$

for $l = 0, 1, \ldots$ The key to solving (4.1) is finding the minimum norm solution to the linear system $\widehat{G}(\theta^l)\eta =$ $\nabla_{\theta} F(\theta^l)$ for η . Minimum norm solutions of a linear system can be done by numerous existing methods, such as MINRES [35]. In these methods, it is instrumental to implement the matrix-vector product with the matrix $\widehat{G}(\theta^l).$

In our experiments, we use deep neural networks as the push-forward map T_{θ} . There are several choices of the neural network architecture. We can use the invertible neural networks (e.g., normalizing flow [34], Real NVP [12] and neural ODE [10]) or non-invertible neural networks (e.g., the multi-layer perceptron or ResNet [17]), both has its own advantages. Normalizing flow and continuous normalizing flow simplify the computation of log determinant of Jacobian matrix of the map, so we can easily compute the density function. For the experiment of the Fokker-Planck equation, we use a 60-layer normalizing flow as the push-forward map T_{θ} . For the experiments of the Porous-medium equation and the aggregation model, we pick the residual neural network as push-forward map:

$$T_{\theta} = Id + R_{\theta}$$

where Id is the identity map and $R_{\theta} : \mathbb{R}^d \to \mathbb{R}^d$ is a standard multilayer perceptron. We take the hyperbolic tangent function as the activation function since we require the second-order derivative in the computation of numerical integrals. The bias for the output layer in f_{θ} is set to be None. To solve the linear system $\widehat{G}(\theta)\dot{\theta} = -\nabla_{\theta}F(\theta)$, we utilized MINRES with tolerance $3 \cdot 10^{-4}$ in our experiments.

To initialize $\theta^0 := \theta(0)$, it depends on the given information of the initial distribution. If the initial probability density function is given and easy to generate samples from, such as Gaussian distribution or Gaussian mixture distribution, then one can initialize the $\theta(0)$ such that $T_{\theta(0)}$ is an identity map. If the initial samples are given but the initial probability density function is unknown, one can still initialize the $\theta(0)$ to make $T_{\theta(0)}$ an identity map. If the given initial density function is hard to sample from, one may use Gaussian distribution as a reference distribution and utilize methods such as MCMC [4] and Wasserstein GAN [2] to enforce the initial distribution.

If the probability density function is desired, one can consult with density estimation methods, such as kernel density estimation, while it's not the focus of this paper. In our experiments, we're given the initial density functions that are easy to sample, hence we initialize the push-forward map as an identity map, i.e., $\rho_{\theta(0)} dx = d(T_{\theta(0)\sharp}\lambda)(\cdot) = d\lambda(\cdot) = \rho(0, \cdot) dx.$

The numerical scheme is summarized in Algorithm 4.1.



Fig. 1: Sample plots of computed ρ_{θ} at different time t for Fokker-Planck equation with the Styblinski-Tang function as the potential function V(x)

Algorithm 4.1 Parameterized Wasserstein gradient flow solver
Initialize the neural network T_{θ}
for $l = 0, \cdots, K - 1$ do
Sample $\{X_1, \cdots, X_N\}$, evaluate $\nabla_{\theta} F(\theta^l)$
Solve η as the minimum norm solution to the linear system $\widehat{G}(\theta^l)\eta = -\nabla_{\theta}F(\theta^l)$
Set $\theta^{l+1} = \theta^l + h\eta$
end for
return θ^K , T_{0K}

5. Numerical examples. In this section, we test the proposed PWGF method, Algorithm 4.1, on three WGFs. Depending on the energy functional \mathcal{F} , we choose push-forward maps as ResNet [17], or normalizing flow [21]. Details are shown below.

5.1. Fokker-Planck equation. We first test Algorithm 4.1 on a 30-dimensional Fokker-Planck equation. We set the coefficient D = 1 and choose the potential function V to be the Styblinski-Tang function in (3.4):

(5.1)
$$V(x) = \frac{3}{50} \Big(\sum_{i=1}^{d} x_i^4 - 16x_i^2 + 5x_i \Big).$$

The initial condition is set to be a Gaussian distribution with mean $\mu_0 = \mathbf{0}$ and variance $\sigma_0 = \mathbf{1}$. We use an *M*-layer normalizing flow (M = 40) as the push-forward map T_{θ} :

(5.2)
$$T_{\theta} = f_M \circ f_{M-1} \circ \cdots \circ f_2 \circ f_1,$$

where f_j $(1 \leq j \leq M)$ is:

$$f_i(x) = x + \tanh(w_i^{\top}x + b_i)u_i$$

Here $w_j, u_j \in \mathbb{R}^d$ and $b_j \in \mathbb{R}$, and $\theta = \{w_j, b_j, u_j : 1 \leq j \leq M\}.$

We run Algorithm 4.1 with time step size h = 0.005. 2,000 samples are generated from ρ_{θ} and their projection of these samples onto the 2-dimensional subspace of (x_1, x_2) are plotted. Notice that V(x) has 4 centers, and the push-forward samples follow a density that captures this structure as shown in Figure 1. The KL divergence (3.3) is evaluated empirically with all samples by

(5.3)
$$\mathcal{H}_{KL}(\rho \| \rho_*) = \int \frac{1}{D} V(x) \rho(x) + \rho(x) \log \rho(x) \, dx + \log Z_D$$
$$\approx \frac{1}{N} \sum_{i=1}^N \frac{1}{D} V(x^{(i)}) + \log \rho(x^{(i)}) + \log Z_D, \quad \text{where } \{x^{(i)}\}_{i=1}^N \sim \rho_i$$

We plot the KL divergence curve following the solution of PWGF in Figure 2, which shows a clear decreasing trend.

To evaluate the improvement in its computation efficiency with the new metric $G(\theta)$, we compare the running time of PWGF Algorithm 4.1 with the PFPE algorithm proposed in [29]. We take the 2-dimensional



Fig. 2: The decay of KL divergence along the solution of PWGF for the Fokker-Planck equation with Styblinski-Tang potential in the 30-dimensional example.

FPE with V(x) defined in (5.1). For both algorithms, we use the same neural network structure (40-layer normalizing flow) and the same sample size (12000) to evaluate the KL divergence. The ψ neural network in PFPE, as well as the training parameters, are set as its default values as reported in [29]. We run both algorithms with the same step size. It takes 14.26 seconds for PWGF to finish 10 time steps, while 6093.58 seconds for PFPE to finish 10 time steps, indicating a more than 400 times speed up. Both codes are run on an NVIDIA RTX-3070 GPU with CUDA enabled.

5.2. Porous medium equation. We apply Algorithm 4.1 to solve the porous medium equation (3.9). In recent years, the porous medium equation has been solved by several other methods, such as physics-informed neural networks (PINN) [37] which computes the density function directly in high dimensions. For all three examples of porous-medium equation, we use a neural network with residual structure as the push-forward map

(5.4)
$$T_{\theta} = Id + R_{\theta}$$

where R_{θ} is a multi-layer perceptron with 3 hidden layers and 100 neurons in each hidden layer.

An example with exact weak solution: Suppose the initial condition $\rho(x,0) = \delta(x)$ is the Dirac delta function.

(5.5)
$$\frac{\partial \rho(x,t)}{\partial t} = \Delta \rho^m, \quad x \in \mathbb{R}^d$$

(5.6)
$$\rho(x,0) = \delta(x)$$

The exact weak solution to (5.5) is

(5.7)
$$\rho(x,t) = t^{-\alpha} F\left(\frac{x}{t^{\beta}}\right), \text{ where } F(\xi) = (C - k\xi^2)_+^{\frac{1}{m-1}},$$

and $(s)_{+} = \max\{s, 0\}$. The constants are

$$\alpha = \frac{d}{d(m-1)+2}, \quad \beta = \frac{\alpha}{d}, \quad k = \frac{(m-1)\alpha}{2md}$$

The constant C > 0 in (5.7) can be uniquely determined by normalizing the mass as

$$\int \rho \, dx = 1,$$

which gives

$$C = \left[\frac{\left(\frac{k}{\pi}\right)^{\frac{d}{2}}\Gamma(\frac{d}{2})}{d \cdot B(\frac{d}{2}, \frac{m}{m-1})}\right]^{\frac{1}{\gamma}},$$



Fig. 3: Sample plots of computed ρ_{θ} at different time t for Porous Medium equation with Dirac Delta function as the initial condition for d = 2 and l = 2.4. The figures are plotted with 5000 samples. In the level curves, darker colors correspond to smaller values to emphasize the support.

where $B(\cdot, \cdot)$ and $\Gamma(\cdot)$ are beta and gamma functions respectively and $\gamma = \frac{d}{2(m-1)\alpha}$ [36].

For the experiments of d = 2, the parameter m is set to 2.4, for the experiment of d = 5, the parameter m is set to 3, and for the experiment of d = 15, the parameter m is set to 2. Since the initial condition is a Dirac delta mass which is impossible to be represented by deep neural networks, we set the initial time to a positive value. More precisely, we compute the dynamics of the samples generated from the probability density ρ from t = 0.1 to t = 0.6 for d = 2 and from t = 1 to t = 1.5 for both d = 5 and d = 15. In all experiments, the step size $h = 10^{-4}$, and the number of time steps is T = 5000. The sample size to evaluate $\hat{G}(\theta)$ and $F(\theta)$ are both 30,000. The graph of the samples with d = 2, d = 5 and d = 15 are shown in Figure 3, Figure 4 and Figure 5 respectively. In the plots, the blue points are the computed samples and the colored curves are the level sets of the exact solutions. We select darker colors representing smaller values when plotting the level curves to clearly show the boundary of the finite support of the exact solution. In the example with d = 2, though the same as the exact solution in the sense that they are compactly supported. In the example with d = 15, all samples are still within the compact support, despite that sampling from a density function in high dimensions remains as a challenging problem. All experiments run on an NVIDIA A100 GPU with 40GB GPU memory.

An Example with Gaussian Mixture Distribution as the Initial Condition: In this example, we apply Algorithm 4.1 to the porous medium equation with d = 10, m = 3 and a Gaussian mixture distribution as the initial condition. The equations are

(5.8)
$$\frac{\partial \rho}{\partial t} = \Delta(\rho^m)$$

(5.9)
$$\rho(0, \cdot) = 0.2 \cdot \mathcal{N}(\mu_1, \sigma_1) + 0.8 \cdot \mathcal{N}(\mu_2, \sigma_2)$$

where $\mu_1 = (0, 0, \dots, 0)^T \in \mathbb{R}^{10}, \mu_2(2, 2, \dots, 2)^T \in \mathbb{R}^{10}, \sigma_1 = \text{diag}(0.1, \dots, 0.1)^T \text{ and } \sigma_2 = \text{diag}(0.2, \dots, 0.2)^T.$



Fig. 4: Sample plots of computed ρ_{θ} at different time t for the porous medium equation with Dirac Delta function as the initial condition for d = 5 and l = 3. The figures are plotted with 5000 samples. In the level curves, darker colors correspond to smaller values to emphasize the support.

The sample size to evaluate $\hat{G}(\theta)$ and $F(\theta)$ are both 15,000. The computed samples are shown in Figure 6 plotted with 5000 samples. The exact solution to this initial value problem remains open but our method can provide a simulated sample dynamics.

An Example with Mixed Gaussian Distribution and Uniform Distribution as the Initial Condition: In this example, we apply Algorithm 4.1 to the porous medium equation with d = 5, m = 3 and a mixture of Gaussian and uniform distributions as the initial condition. The equations are

(5.10)
$$\frac{\partial \rho}{\partial t} = \Delta(\rho^m)$$

(5.11)
$$\rho(0,\cdot) = 0.2 \cdot \mathcal{N}(\mu,\sigma) + 0.8 \cdot \mathcal{U}_5([-1,1]^5)$$

where $\mu = (2, 2, \dots, 2)^T \in \mathbb{R}^5$, $\sigma = (0.1, 0.1, \dots, 0.1)^T \in \mathbb{R}^5$ and $\mathcal{U}_5([-1, 1]^5)$ is a uniform distribution in $[-1, 1]^5$. Note that this equation doesn't have a closed-form solution and it's not tractable in high dimensions. The sample size to evaluate $\widehat{G}(\theta)$ and $F(\theta)$ are both 20,000. The computed samples are shown in Figure 7 plotted with 5000 samples.

5.3. Aggregation model as a Wasserstein gradient flow. In this example, we apply Algorithm 4.1 to the aggregation problem with d = 2 and a = 4, b = 2, i.e., the energy functional \mathcal{F} is set as in (3.10) with J(x) given by

(5.12)
$$J(x) = \frac{|x|^4}{4} - \frac{|x|^2}{2}.$$



Fig. 5: Sample plots of computed ρ_{θ} at different time t for porous medium equation with Dirac Delta function as the initial condition for d = 15. The figures are plotted with 5000 samples. In the level curves, darker colors correspond to smaller values to emphasize the support.



Fig. 6: Sample plots of computed ρ_{θ} at different time t for porous medium equation with Gaussian mixture distribution as the initial condition. The figures are plotted with 5000 samples.

We set the initial condition to be Gaussian distribution with mean $\mu_0 = (1.25, 1.25)^{\top}$ and variance γI where $\gamma = 0.6$.

(5.13)
$$\rho_0(x) = \frac{1}{\sqrt{2\pi\gamma}} e^{-\frac{|x-\mu_0|^2}{2\gamma^2}}$$



Fig. 7: Sample plots of computed ρ_{θ} at different time t for porous medium equation with the mixture of Gaussian distribution and uniform distribution as the initial condition. The graphs are plotted with 5000 samples.



Fig. 8: Sample plots of computed ρ_{θ} at different time t for aggregation problem

We use a neural network with residual structure as the push-forward map in this example:

$$(5.14) T_{\theta} = Id + R_{\theta}$$

where R_{θ} is multi-layer perceptron with 2 hidden layers and 50 neurons in each hidden layer. The sample size to evaluate $\hat{G}(\theta)$ and $F(\theta)$ are both 10,000.

The steady solution ρ_* is a Dirac distribution uniformly concentrated on the ring with radius 0.5 centered at μ_0 . ρ converges to ρ_* when $t \to \infty$. This evolution is captured as shown in Figure 8, which shows the distribution $\rho_{\theta(t)}$ of samples from at different times t. Note that, despite of the strong singularity of ρ_* , the samples generated by Algorithm 4.1 can approximate this distribution well.

6. Discussion. We proposed a new numerical approach to solve Wasserstein gradient flows (WGFs), which is particularly scalable for high-dimensional cases. Our approach used general reduced-order models, like deep neural networks, to parameterize the push-forward maps such that they can push a simple reference density to the density solving the given WGF. Essentially, the gradient flow defined on the infinite-dimensional Wasserstein manifold is reduced to a finite-dimensional dynamical system, the PWGF, for the parameters used in the reduced-order model. At the core of our design is the pullback Wasserstein metric on the parameter space. It facilitates the derivation, simplifies the numerical computation, and plays a pivotal role in the error-bound estimates that offer theoretical assurance for PWGF to the original WGF.

Though the parametrization of the WGF provides a fast algorithm, the accuracy and speed of the algorithm largely rely on the linear system solver for the neural network parameter θ . Since the metric matrix $\hat{G}(\theta)$ is semi-positive definite and can be degenerate in some scenarios, the minimal residual method is utilized to find the least square solution. Our method is sample-based and provides the value of the density function at sampled points, it has the potential to adapt to tasks such as developing a fast sampler of target distribution by evolving samples from the initial probability density following the gradient flow or image generation following certain dynamics of probability densities. In these scenarios, as the task is more complex, the deep neural network for parametrizing the push-forward map could be large or have sophisticated architecture. One may consider dimension reduction techniques to reduce the dimension of the parameter space. In addition, other improvements such as more efficient linear system solvers or preconditioning techniques may also be considered to boost the accuracy and speed. Meanwhile, better understanding on δ_0 in the error analysis needs to be analyzed regarding various neural network structures.

Acknowledgments. This research is partially supported by NSF grants DMS-1925263, DMS-2152960, DMS-2307465, DMS-2307466, and ONR grant N00014-21-1-2891. All the authors made equal contributions.

References.

- [1] W. ANDERSON AND M. FARAZMAND, Evolution of nonlinear reduced-order solutions for PDEs with conserved quantities, SIAM Journal on Scientific Computing, 44 (2022), pp. A176–A197.
- [2] M. ARJOVSKY, S. CHINTALA, AND L. BOTTOU, Wasserstein generative adversarial networks, in International conference on machine learning, PMLR, 2017, pp. 214–223.
- [3] J.-D. BENAMOU, G. CARLIER, Q. MÉRIGOT, AND E. OUDET, Discretization of functionals involving the monge-ampère operator, Numerische mathematik, 134 (2016), pp. 611–636.
- [4] B. A. BERG, Markov chain Monte Carlo simulations and their statistical analysis: with web-based Fortran code, World Scientific Publishing Company, 2004.
- [5] A. BLANCHET AND P. LAURENÇOT, The parabolic-parabolic keller-segel system with critical diffusion as a gradient flow in \mathbb{R} d, $d \ge 3$, Communications in Partial Differential Equations, 38 (2013), pp. 658–686.
- [6] S. C. BRENNER, The mathematical theory of finite element methods, Springer, 2008.
- [7] J. BRUNA, B. PEHERSTORFER, AND E. VANDEN-EIJNDEN, Neural Galerkin schemes with active learning for high-dimensional evolution equations, Journal of Computational Physics, 496 (2024), p. 112588.
- [8] G. CARLIER, V. DUVAL, G. PEYRÉ, AND B. SCHMITZER, Convergence of entropic schemes for optimal transport and gradient flows, SIAM Journal on Mathematical Analysis, 49 (2017), pp. 1385–1418.
- [9] J. A. CARRILLO, K. CRAIG, L. WANG, AND C. WEI, Primal dual methods for wasserstein gradient flows, Foundations of Computational Mathematics, (2022), pp. 1–55.
- [10] R. T. CHEN, Y. RUBANOVA, J. BETTENCOURT, AND D. K. DUVENAUD, Neural ordinary differential equations, Advances in neural information processing systems, 31 (2018).
- [11] S. CHEWI, T. LE GOUIC, C. LU, T. MAUNU, AND P. RIGOLLET, Svgd as a kernelized wasserstein gradient flow of the chi-squared divergence, Advances in Neural Information Processing Systems, 33 (2020), pp. 2098–2109.
- [12] L. DINH, J. SOHL-DICKSTEIN, AND S. BENGIO, *Density estimation using real nvp*, arXiv preprint arXiv:1605.08803, (2016).
- [13] Y. DU AND T. A. ZAKI, Evolutional deep neural network, Physical Review E, 104 (2021), p. 045303.
- [14] J. FAN, Q. ZHANG, A. TAGHVAEI, AND Y. CHEN, Variational wasserstein gradient flow, in International Conference on Machine Learning, PMLR, 2022, pp. 6185–6215.
- [15] R. C. FETECAU, Y. HUANG, AND T. KOLOKOLNIKOV, Swarm dynamics and equilibria for a nonlocal aggregation model, Nonlinearity, 24 (2011), p. 2681.
- [16] N. GABY, X. YE, AND H. ZHOU, Neural control of parametric solutions for high-dimensional evolution pdes, SIAM Journal on Scientific Computing, 46 (2024), pp. C155–C185.
- [17] K. HE, X. ZHANG, S. REN, AND J. SUN, Deep residual learning for image recognition, in Proceedings of the IEEE conference on computer vision and pattern recognition, 2016, pp. 770–778.
- [18] R. HOLLEY AND D. W. STROOCK, Logarithmic sobolev inequalities and stochastic ising models, Journal of Statistical Physics, 46 (1987), pp. 1159–1194, https://doi.org/10.1007/BF01011161.
- [19] Z. HU, C. LIU, Y. WANG, AND Z. XU, Energetic variational neural network discretizations to gradient flows, arXiv preprint arXiv:2206.07303, (2022).
- [20] R. JORDAN, D. KINDERLEHRER, AND F. OTTO, The variational formulation of the fokker-planck equation, SIAM journal on mathematical analysis, 29 (1998), pp. 1–17.
- [21] I. KOBYZEV, S. J. PRINCE, AND M. A. BRUBAKER, Normalizing flows: An introduction and review of current methods, IEEE transactions on pattern analysis and machine intelligence, 43 (2020), pp. 3964– 3979.
- [22] A. J. LADD AND P. SZYMCZAK, Reactive flows in porous media: challenges in theoretical and numerical

methods, Annual review of chemical and biomolecular engineering, 12 (2021), pp. 543–571.

- [23] J. D. LAFFERTY, The Density Manifold and Configuration Space Quantization, Transactions of the American Mathematical Society, 305 (1988), pp. 699–741.
- [24] W. LEE, L. WANG, AND W. LI, Deep JKO: time-implicit particle methods for general nonlinear gradient flows, arXiv preprint arXiv:2311.06700, (2023).
- [25] R. J. LEVEQUE, Finite difference methods for ordinary and partial differential equations: steady-state and time-dependent problems, SIAM, 2007.
- [26] W. LI, S. LIU, H. ZHA, AND H. ZHOU, Parametric Fokker-Planck Equation, Springer International Publishing, 2019, p. 715–724, https://doi.org/10.1007/978-3-030-26980-7_74, http://dx.doi.org/10.1007/ 978-3-030-26980-7_74.
- [27] W. LI, J. LU, AND L. WANG, Fisher information regularization schemes for wasserstein gradient flows, Journal of Computational Physics, 416 (2020), p. 109449.
- [28] W. LI AND J. ZHAO, Wasserstein information matrix, Information Geometry, (2023), pp. 1–53.
- [29] S. LIU, W. LI, H. ZHA, AND H. ZHOU, Neural parametric fokker-planck equation, SIAM Journal on Numerical Analysis, 60 (2022), pp. 1385–1449.
- [30] P. MOKROV, A. KOROTIN, L. LI, A. GENEVAY, J. M. SOLOMON, AND E. BURNAEV, Large-scale wasserstein gradient flows, Advances in Neural Information Processing Systems, 34 (2021), pp. 15243– 15256.
- [31] L. NURBEKYAN, W. LEI, AND Y. YANG, Efficient natural gradient descent methods for large-scale pdebased optimization problems, SIAM Journal on Scientific Computing, 45 (2023), pp. A1621–A1655.
- [32] F. OTTO, The Geometry of Dissipative Evolution Equations: The Porous Medium Equation, Communications in Partial Differential Equations, 26 (2001), pp. 101–174.
- [33] G. PEYRÉ, Entropic approximation of wasserstein gradient flows, SIAM Journal on Imaging Sciences, 8 (2015), pp. 2323–2351.
- [34] D. REZENDE AND S. MOHAMED, Variational inference with normalizing flows, in International conference on machine learning, PMLR, 2015, pp. 1530–1538.
- [35] Y. SAAD, Iterative methods for sparse linear systems, SIAM, 2003.
- [36] J. L. VÁZQUEZ, The porous medium equation: mathematical theory, Oxford University Press, 2007.
- [37] M. WANG AND J. LU, Neural network-based variational methods for solving quadratic porous medium equations in high dimensions, Communications in Mathematics and Statistics, 11 (2023), pp. 21–57.
- [38] H. WU, Theory and computation of Wasserstein geometric flows with application to time-dependent Schrodinger equation, (2023). Publisher: Georgia Institute of Technology.
- [39] H. WU, S. LIU, X. YE, AND H. ZHOU, *Parameterized wasserstein hamiltonian flow*, arXiv preprint arXiv:2306.00191, (2023).
- [40] M. YI AND S. LIU, Bridging the gap between variational inference and wasserstein gradient flows, 2023, https://arxiv.org/abs/2310.20090.
- [41] X. ZUO, J. ZHAO, S. LIU, S. OSHER, AND W. LI, Numerical analysis on neural network projected schemes for approximating one dimensional wasserstein gradient flows, 2024, https://arxiv.org/abs/2402.16821.