No ν s is Good News

Nathaniel Craig^{1,2}, Daniel Green³, Joel Meyers⁴, and Surjeet Rajendran⁵

¹Department of Physics, University of California, Santa Barbara, CA 93106, USA

²Kavli Institute for Theoretical Physics, Santa Barbara, CA 93106, USA

³Department of Physics, University of California, San Diego, La Jolla, CA 92093, USA

⁴Department of Physics, Southern Methodist University, Dallas, TX 75275, USA

⁵Department of Physics & Astronomy, The Johns Hopkins University,

Baltimore, MD 21218, USA

Abstract

The baryon acoustic oscillation (BAO) analysis from the first year of data from the Dark Energy Spectroscopic Instrument (DESI), when combined with data from the cosmic microwave background (CMB), has placed an upper-limit on the sum of neutrino masses, $\sum m_{\nu} < 70 \text{ meV} (95\%)$. In addition to excluding the minimum sum associated with the inverted hierarchy, the posterior is peaked at $\sum m_{\nu} = 0$ and is close to excluding even the minumum sum, 58 meV at 2σ . In this paper, we explore the implications of this data for cosmology and particle physics. The sum of neutrino mass is determined in cosmology from the suppression of clustering in the late universe. Allowing the clustering to be enhanced, we extended the DESI analysis to $\sum m_{\nu} < 0$ and find $\sum m_{\nu} = -160 \pm 90$ meV (68%), and that the suppression of power from the minimum sum of neutrino masses is excluded at 99% confidence. We show this preference for negative masses makes it challenging to explain the result by a shift of cosmic parameters, such as the optical depth or matter density. We then show how a result of $\sum m_{\nu} = 0$ could arise from new physics in the neutrino sector, including decay, cooling, and/or time-dependent masses. These models are consistent with current observations but imply new physics that is accessible in a wide range of experiments. In addition, we discuss how an apparent signal with $\sum m_{\nu} < 0$ can arise from new long range forces in the dark sector or from a primordial trispectrum that resembles the signal of CMB lensing.

Contents

1	Intr	oduction	3		
2	Neutrino Mass and DESI				
	2.1	How Neutrino Mass is Measured	5		
	2.2	Negative Neutrino Mass	7		
	2.3	Influence of Cosmic Parameters	9		
3	Vanishing Neutrino Mass				
	3.1	Decays	16		
	3.2	Annihilation	17		
	3.3	Cooling and Heating	18		
	3.4	Time Varying Mass	21		
	3.5	Mirror Sectors and Relation to the Hubble Tension	21		
4	Negative "Neutrino Mass"				
	4.1	Dark Matter with Long Range Forces	22		
	4.2	Primordial Trispectrum	24		
5	Conclusions				
A	A The Suppression of Clustering				
Re	References				

1 Introduction

The cosmological measurement of the sum of neutrino masses, $\sum m_{\nu}$, is one of the most anticipated results from the coming generation of cosmic surveys [1–3]. From the measurement of neutrino flavor oscillations [4], which precisely determine the mass-squared splittings between neutrino mass eigenstates, it can be inferred that the sum of neutrino masses is necessarily greater than 58 meV. This provides a concrete prediction within the standard cosmological model that should be measurable (or detectable) with planned observations [5,6].

The Dark Energy Spectroscopic Instrument (DESI) [7] is expected to provide the necessary increase in sensitivity to $\sum m_{\nu}$ to measure the minimum sum at 2 to 3σ [5, 6]. Cosmological measurements of neutrino mass rely on the measurement of the clustering of matter on scales smaller than the free-streaming length of neutrinos [1]. A universe containing massive neutrinos will exhibit suppressed matter clustering compared to a universe with only massless neutrinos. This measurement can be achieved by combining observations of the cosmic microwave background (CMB) with the measurement of the baryon acoustic oscillations (BAO). The amplitude of clustering can be determined from the measurement of the CMB lensing power spectrum, and this amplitude is compared to what would be expected in a universe with only massless neutrinos [8]. In the absence of massive neutrinos, the amplitude of matter clustering is determined by the matter density and the primordial amplitude of scalar fluctuations. Measurements of the CMB angular power spectra allow for a determination of the primordial fluctuation amplitude. BAO measurements are needed to measure the abundance of non-relativistic matter to sufficient accuracy to isolate the effect neutrino mass [9].

The release of the first year BAO analysis with DESI [10], combined with data from the CMB (Planck 2018 [11,12] and ACT DR6 lensing [13,14]), showed a remarkable upper-limit on $\sum m_{\nu}$, reaching

$$\sum m_{\nu} < 70 \,\mathrm{meV} \,(95\%) \ . \tag{1.1}$$

This is consistent with an earlier constraint from (e)BOSS of $\sum m_{\nu} < 82 \text{ meV}$ [15] using CMB+BAO+Shape parameters (see also [16]). The DESI result is sufficient to exclude the minimum mass for an inverted neutrino mass hierarchy, 100 meV, at ~ 3σ . However, what is also noteworthy is that the posterior peaks at $\sum m_{\nu} = 0$ and is very close to putting 58 meV in tension with observations.

In this paper, we will explore the current constraints on $\sum m_{\nu}$ and what an exclusion of $\sum m_{\nu} = 58$ meV would mean for cosmology and particle physics. First, we will examine the current measurement and how it depends on different types of surveys. One particularly noteworthy aspect of the DESI measurement is that it appears to favor $\sum m_{\nu} < 0$, though that region of parameter space was excluded from the DESI analysis by imposing a prior that $\sum m_{\nu}$ is positive. Although negative neutrino masses are unphysical, a preference in the data for $\sum m_{\nu} < 0$ may simply reflect an excess of clustering in the late universe,



Figure 1: Posterior of neutrino mass in eV inferred from Planck + ACT Lensing + DESI data. The blue line shows constraints on a model with a physical neutrino mass, the orange line shows constraints where the neutrino mass is parametrized as an effect on the CMB lensing power spectrum and restricted to be positive, and the green line shows constraints on a parametrized neutrino mass that is allowed to be negative. The best fit for the parametrized neutrino mass is $\sum \tilde{m}_{\nu} = -160$ meV, and the minimal neutrino mass of 58 meV is disfavored at 3σ . For details about the parametrization of negative neutrino mass and the data sets used, see Section 2.2.

rather than a deficit caused by free streaming neutrinos. We use this idea to define a neutrino mass, $\sum \tilde{m}_{\nu}$, that is allowed to be negative and perform the same analysis as DESI without the positive mass prior. We find that that data does prefer negative mass, $\sum \tilde{m}_{\nu} = -160 \pm 90 \text{ meV} (68\%)$, and corresponds to a 3σ exclusion of the minimum neutrino mass. The full posterior is shown in Figure 1.

The preference of the current measurement for negative $\sum m_{\nu}$ is particularly important as it affects the bias in the measurement of cosmic parameters, particularly the optical depth, τ , that would be required to explain the current limits. For $\ell > 30$, the CMB is only sensitive to the combination $A_s e^{-2\tau}$, where A_s is the amplitude of primordial scalar fluctuations. The determination of τ is therefore essential for determining A_s and suppression of power a late times, but requires (challenging) large angular scale measurements of the CMB. It is plausible that $\sum m_{\nu} = 0$ could be explained by a statistical or systematic shift in τ , but it is far more challenging to explain $\sum m_{\nu} = -160$ meV in this way.

An absence of the neutrino mass signal, while forbidden in the Standard Model (plus neutrino masses), could be a natural consequence of a wide variety of beyond the Standard Model (BSM) scenarios. The most straightforward mechanisms to eliminate the signal would be to eliminate the SM neutrinos via decay (or annihilation), cool the neutrinos so that they behave like dark matter, or change their mass over cosmological history. Simple models for all three scenarios can be derived from new interactions in the neutrino sector that are weakly constrained by experiments. On the other hand, the CMB does provide stringent constraints on the parameter space of these models, as measurements of N_{eff} are in good agreement with the expected temperature [17] and free-streaming [18–20] of the cosmic neutrino background (C ν B). Nevertheless, there have been hints of neutrino interactions [21–25] in cosmic data that may also point to new physics of this kind.

Negative neutrino masses, $\sum m_{\nu} < 0$, are representative of enhanced clustering of matter, rather than any physical property of the neutrinos themselves. This kind of enhanced clustering can be achieved by changing the long range forces that act on matter. We discuss one simple mechanism, which is to introduce a new scalar force that acts only on the dark matter. Such forces are more weakly constrained than fifth forces acting on SM particles and thus could explain our signal without being in tension with other constraints. Alternatively, a CMB lensing measurement with $\sum m_{\nu} < 0$ points to a larger than expected CMB trispectrum, which could result from a non-zero primordial trispectrum. These scenarios will all be testable with current and/or future cosmic data.

This paper is organized as follows: In Section 2, we review the measurement of $\sum m_{\nu}$ and extend the analysis to negative masses. We discuss what shifts in cosmic data would be required to make these measurements consistent with conventional neutrino physics. In Section 3, we present models that could explain $\sum m_{\nu} = 0$ with new physics in the neutrino sector. In Section 4, we present models that could explain a cosmological inference of negative neutrino masses. We conclude in Section 5. Appendix A, we review the physics origin of the suppression of structure due to massive neutrinos.

2 Neutrino Mass and DESI

2.1 How Neutrino Mass is Measured

In order to understand what an apparent measurement of $\sum m_{\nu} = 0$ would mean, we first need to review exactly what measurements allow us to infer $\sum m_{\nu}$ (see also [26, 27] for review). We will assume that $\sum m_{\nu} \approx 60$ meV, as this is the minimum sum consistent with neutrino oscillation experiments and is therefore the minimum value that would need to be excluded in order to favor $\sum m_{\nu} = 0$.

Cosmic neutrinos are relativistic in the early universe, but become non-relativistic when their propagation speed, c_{ν} , drops well below the speed of light. In a ACDM + m_{ν} cosmology, the typical neutrino speed is given by

$$c_{\nu} = \frac{\langle p_{\nu} \rangle}{m_{\nu}} = \frac{3T_{\nu}}{m_{\nu}} \approx 1.0 \times 10^{-2} \left(\frac{50 \,\mathrm{meV}}{m_{\nu}}\right) \,(1+z) \,,$$
 (2.1)

where we have set c = 1. As a result, the redshift where the heaviest neutrino becomes non-relativistic is $z_{\nu} \approx 100$. For $z < z_{\nu}$, the energy density of neutrinos redshifts like non-relativistic matter so that

$$\Omega_m = \Omega_c + \Omega_b + \Omega_\nu \ . \tag{2.2}$$

However, the neutrinos are still sufficiently hot that they do not cluster on scales below their effective Jeans scale. In terms of wavenumber, this free-streaming scale is given by

$$k_{\rm fs} = \sqrt{\frac{3}{2}} \frac{aH}{c_{\nu}} = 0.04 \, h \, {\rm Mpc}^{-1} \times \frac{1}{1+z} \left(\frac{\sum m_{\nu}}{58 \, {\rm meV}}\right) \,. \tag{2.3}$$

Because neutrinos don't cluster, the amplitude of clustering of matter, defined by the matter power spectrum

$$P(k) = \langle \delta_m(\vec{k}) \delta_m(\vec{k}') \rangle' , \qquad (2.4)$$

is suppressed on scales smaller than the neutrino free-streaming scale $k \gg k_{\rm fs}$

$$P^{(\sum m_{\nu})}(k \gg k_{\rm fs}, z) \approx \left(1 - 2f_{\nu} - \frac{6}{5}f_{\nu}\log\frac{1+z_{\nu}}{1+z}\right)P^{(\sum m_{\nu}=0)}(k \gg k_{\rm fs}, z) , \qquad (2.5)$$

where $f_{\nu} = \Omega_{\nu}/\Omega_m$ is the fraction of non-relativistic matter in the form of neutrinos, $\delta_m \equiv \delta \rho_m/\bar{\rho}_m$ is the density contrast of non-relativistic matter, and the prime on the correlation function means that the delta function has been omitted. The suppression in this formula is the result of two distinct physical effects (see Appendix A for a derivation). The first term, $-2f_{\nu}$, reflects the reduced fraction of matter that is actually clustering. The second, $-\frac{6}{5}f_{\nu}\log\frac{1+z_{\nu}}{1+z}$, is due to reduced rate of growth of the dark matter perturbations in the presence of matter that doesn't cluster. Using

$$\Omega_{\nu}h^{2} = 6 \times 10^{-4} \left(\frac{\sum m_{\nu}}{58 \,\mathrm{meV}}\right) \to f_{\nu} \approx 4 \times 10^{-3} , \qquad (2.6)$$

the suppression of the matter power spectrum at z = 1 is expected to be

$$P^{(\sum m_{\nu}=58\,\mathrm{meV})}(k\gg k_{\mathrm{fs}},z)\approx (1-0.02)P^{(\sum m_{\nu}=0)}(k\gg k_{\mathrm{fs}},z) \ . \tag{2.7}$$

Therefore, the signal we are looking for is a 2% suppression of power on small scales around $z = \mathcal{O}(1)$.

Galaxy surveys like DESI do not directly measure P(k) and instead primarily measure the clustering of galaxies. The power spectrum of galaxy overdensity has an overall amplitude that depends on the details of galaxy formation, and the baryonic physics inherent in galaxy formation is understood with insufficient precision to directly extract the amplitude of P(k) from these measurements. The best current measurements of the matter power spectrum come from gravitational lensing of the CMB. The CMB lensing convergence power spectrum $C_{\ell}^{\kappa\kappa}$ is given in the Limber approximation by [28]

$$C_{\ell}^{\kappa\kappa} \approx 2\pi^2 \ell \int_{\eta_*}^{\eta_0} \eta \mathrm{d}\eta \mathcal{P}_{\Psi} \left(\ell / (\eta_0 - \eta); \eta \right) \left(\frac{\eta_* - \eta}{(\eta_0 - \eta_*)(\eta_0 - \eta)} \right)^2 , \qquad (2.8)$$

where η is the conformal time with η_{\star} and η_0 denoting the times of recombination and z = 0 respectively. We also defined \mathcal{P}_{Ψ} as power spectrum of the Weyl potential, Ψ , which can be written in terms of the matter power spectrum as

$$\mathcal{P}_{\Psi}(k;\eta) = \frac{9\Omega_m^2(\eta)H^4(\eta)}{8\pi^2} \frac{P(k;\eta)}{k} \,. \tag{2.9}$$

Using the fact that the matter power spectrum is proportional to the primordial scalar amplitude A_s , we see that the amplitude of the CMB lensing power spectrum scales as

$$C_{\ell}^{\kappa\kappa} \propto (\Omega_m h^2)^2 A_s \left(1 - 0.02 \frac{f_{\nu}}{4 \times 10^{-3}} \right)$$
 (2.10)

Therefore, in order to measure a three-percent suppression of the lensing power spectrum, we must determine the physical matter density $\Omega_m h^2$ (where $h = H_0/(100 \text{ km s}^{-1} \text{ Mpc}^{-1})$ is the dimensionless Hubble constant) and the primordial scalar amplitude A_s to much better than three-percent accuracy.

The main impact of DESI on the cosmological neutrino mass constraint is to provide a precise measurement of $\omega_m \equiv \Omega_m h^2$ through the constraint on the expansion history from BAO. The impact of changing ω_m on the CMB lensing power spectrum is shown in Figure 2 (for $\sum m_{\nu} = 0$ and compared to the change from introducing $\sum m_{\nu} > 0$). The reduction of ω_m by 1.7% is roughly equivalent to introducing $\sum m_{\nu} = 58$ meV, which implies that a 2σ measurement of the minimum sum requires roughly 0.8% precision in the measurement of ω_m .

2.2 Negative Neutrino Mass

The physical sum of neutrino masses is of course restricted to be positive. However, the combination of cosmological observables that we use to infer the mass of neutrinos are not restricted in this manner. We show in this subsection that the CMB+DESI data in fact prefer a negative neutrino mass (already hinted at in eBOSS [29]), corresponding to increased matter clustering compared to a model with only massless neutrinos.

In order to measure the preference of cosmological data for negative neutrino mass, we require an implementation of the effects of neutrino mass that is allowed to take either sign. The Boltzmann codes CAMB [30, 31] and CLASS [32] model neutrino mass in a way that is subject to the physicality constraint $\sum m_{\nu} > 0$. We modified CAMB to include a new parameter, $\sum \tilde{m}_{\nu}$, which is designed to mimic the effects of neutrino mass, but which



Figure 2: Comparison of the fractional change to the CMB lensing power spectrum from changes to $\Omega_m h^2$ and the introduction of a non-zero neutrino mass.

is not restricted to be positive. Our new parameter simply scales the CMB lensing power spectrum in the same manner that would be expected from $\sum m_{\nu}$. Specifically, we determine the fractional change $A_{\ell}(\sum \tilde{m}_{\nu}) \equiv C_{\ell}^{\kappa\kappa} [\sum m_{\nu}]/C_{\ell}^{\kappa\kappa} [\sum m_{\nu} = 0]$ at fixed values of H_0, ω_m , and ω_b . Once calibrated on positive values of neutrino mass, the effects of $\sum \tilde{m}_{\nu}$ can then be straightforwardly calculated for negative values as well. In the $\Lambda \text{CDM} + \sum \tilde{m}_{\nu}$ cosmology, observables are computed with the physical $\sum m_{\nu} = 0$ and the CMB lensing power spectrum is computed as $C_{\ell}^{\kappa\kappa} = A_{\ell}(\sum \tilde{m}_{\nu})C_{\ell}^{\kappa\kappa}[\sum m_{\nu} = 0]$. The temperature and polarization CMB power spectra are lensed using this modified CMB lensing power spectrum such that the set C_{ℓ}^{TT} , C_{ℓ}^{TE} , C_{ℓ}^{EE} , $C_{\ell}^{\kappa\kappa}$ is calculated self-consistently for each point in parameter space.

This prescription is very similar, though not identical, to the effects of the physical neutrino mass in the regime $\sum m_{\nu} > 0$. In particular, the physical neutrino mass in the $\Lambda \text{CDM} + \sum m_{\nu}$ cosmology contributes to the non-relativistic matter density today $\Omega_m =$ $\Omega_b + \Omega_c + \Omega_{\nu}$. In our $\Lambda \text{CDM} + \sum \tilde{m}_{\nu}$ cosmology, there is no neutrino contribution to Ω_m . As a result, we anticipate that $\sum \tilde{m}_{\nu}$ should exhibit slightly weaker constraints than the physical $\sum m_{\nu}$ when measured using the same data combination. To check this, we derive constraints on three cosmological models: a model with a physical neutrino mass $\Lambda \text{CDM} + \sum m_{\nu}$, a model with our parametrized neutrino mass restricted to positive values $\Lambda \text{CDM} + (\sum \tilde{m}_{\nu} > 0)$, and finally a model with our parametrized neutrino mass with no restriction on sign $\Lambda CDM + \sum \tilde{m}_{\nu}$. We analyze each model using the same data combination.

Boltzmann calculations were carried out using our modified version of CAMB [30, 31]. We utilized the likelihood for CMB temperature and polarization from Planck's 2018 data release [11], along with the combination of ACT DR6 [13,14] and Planck CMB lensing [12], and DESI BAO [10,33,34]. This combination of data is the same as that used by the DESI team to derive cosmological constraints [10]. Our analysis was performed with cobaya [35], using the Markov chain Monte Carlo sampler adapted from CosmoMC [36,37] using the fast-dragging procedure [38]. Analyses were run until the Gelman-Rubin statistic was R - 1 < 0.01.

The results are presented in Table 1 and Figure 3. Notice that the parameter constraints in the $\Lambda \text{CDM} + \sum m_{\nu}$ and $\Lambda \text{CDM} + (\sum \tilde{m}_{\nu} > 0)$ models are nearly identical, showing only slightly weaker constraints on $\sum \tilde{m}_{\nu}$ as compared to the physical $\sum m_{\nu}$. This excellent agreement justifies our prescription for modeling the effects of neutrino mass, with the slightly weaker constraints on $\sum \tilde{m}_{\nu}$ expected from the differing treatment of Ω_m in the two models. Notice that in the $\Lambda \text{CDM} + \sum \tilde{m}_{\nu}$ model, the best-fit value for $\sum \tilde{m}_{\nu}$ is -160 meV, showing a preference for negative neutrino mass, and disfavoring even the minimal sum of neutrino masses inferred from flavor oscillation experiments at 3σ .

We also note in passing that in the $\Lambda \text{CDM} + \sum \tilde{m}_{\nu}$ model, the best-fit value for $S_8 \equiv \sigma_8 (\Omega_m/0.3)^{0.5}$ is lower than in $\Lambda \text{CDM} + \sum m_{\nu}$ by about 1.5σ and has 40% larger error bars (and is also smaller than the value inferred with Planck in the ΛCDM model, for which $S_8 = 0.830 \pm 0.013$ [17]), representing a somewhat smaller S_8 tension [39] when neutrino mass is allowed to be negative.

2.3 Influence of Cosmic Parameters

Optical Depth

The measurement of A_s is limited by our understanding of the optical depth to reionization, τ . Thomson scattering of CMB photons into and out of the line of site by free electrons present after reionization suppresses the amplitude of CMB fluctuations. The observed amplitude of the CMB power spectrum is thereby reduced on small angular scales. CMB observations primarily constrain the combination [17]

$$A_s e^{-2\tau} = (1.884 \pm 0.011) \times 10^{-9} . \tag{2.11}$$

This should be contrasted with the much less precise measurement of the primordial amplitude [17]

$$A_s = (2.100 \pm 0.030) \times 10^{-9} . \tag{2.12}$$

	$\Lambda ext{CDM} + \sum m_ u$	$\Lambda ext{CDM} + (\sum ilde{m}_ u > 0)$	$\Lambda ext{CDM} + \sum ilde{m}_{ u}$
Parameter	68% limits	68% limits	68% limits
$\log(10^{10}A_{ m s})$	3.051 ± 0.014	$3.053\substack{+0.013\\-0.014}$	3.030 ± 0.017
$n_{ m s}$	0.9692 ± 0.0037	0.9686 ± 0.0036	0.9708 ± 0.0038
$100 heta_{ m MC}$	1.04112 ± 0.00029	1.04111 ± 0.00029	1.04118 ± 0.00029
$\Omega_{ m b}h^2$	0.02249 ± 0.00013	0.02248 ± 0.00013	0.02255 ± 0.00014
$\Omega_{ m c}h^2$	0.11852 ± 0.00088	0.11880 ± 0.00088	0.11780 ± 0.00097
$\sum m_ u \left(\sum ilde{m}_ u ight)$	$< 0.0741 \ (95\% \ {\rm CL})$	$< 0.0926 \ (95\% \ {\rm CL})$	$-0.156^{+0.093}_{-0.085}$
$ au_{ m reio}$	0.0585 ± 0.0074	$0.0588^{+0.0069}_{-0.0077}$	0.0510 ± 0.0083
H_0	68.33 ± 0.43	68.43 ± 0.40	68.87 ± 0.45
$\Omega_{ m m}h^2$	0.14131 ± 0.00084	0.14127 ± 0.00083	0.14036 ± 0.00092
σ_8	$0.8175\substack{+0.0076\\-0.0058}$	$0.8246^{+0.0053}_{-0.0060}$	0.8123 ± 0.0078
$S_8 \equiv \sigma_8 (\Omega_m/0.3)^{0.5}$	0.8212 ± 0.0096	0.827 ± 0.010	0.807 ± 0.013

Table 1: Parameter constraints from Planck + ACT lensing + DESI BAO in the three models described in the text. All constraints are given as 68% limits, except for the upper limits on the neutrino mass when it is restricted to be positive, which are reported as 95% CL. Values of neutrino mass are reported in eV and H_0 in km s⁻¹ Mpc⁻¹. In the Λ CDM+ $\sum \tilde{m}_{\nu}$ model, the data favors a negative neutrino mass and disfavors the minimal physical neutrino mass of 58 meV at 3σ .

Noting that for these same analyses,

$$\tau = 0.0544 \pm 0.0073 \,, \tag{2.13}$$

the error on A_s can be directly attributed to the error in τ and not the error in the measurement of $A_s e^{-2\tau}$.

Of all the cosmological parameters defining Λ CDM, the optical depth is the most challenging to measure. For $\ell > 30$, its effects on the CMB are completely degenerate with A_s . It is only on large angular scales that the optical depth leaves a unique imprint, through the production of CMB polarization and the associated 'reionization bump' in the polarization power spectrum. The history of these measurements, shown in Figure 4 has involved significant changes in the central value with relatively small changes in sensitivity.

It is natural to wonder if the apparent measurement of $\sum m_{\nu} = -160$ meV could also be attributed to an error in the measurement of τ . For this to be possible, we would need the true value of A_s to be roughly 8.8% larger, so that the current measurement of the lensing includes the expected suppression of P(k) relative to A_s . This would require a



Figure 3: Triangle plot showing parameter constraints from Planck + ACT lensing + DESI BAO in three models described in the text and shown in Table 1. For the purposes of this plot, we treat the physical neutrino mass and our parametrized version on the same footing. Dashed lines show vanishing neutrino mass $\sum m_{\nu} = 0$ and the minimal sum of neutrino mass $\sum m_{\nu} = 58$ meV. Values of neutrino mass are reported in eV and H_0 in km s⁻¹ Mpc⁻¹.

value of the optical depth larger than that inferred from Planck $\tau = \tau_{\text{Planck}} + \delta \tau$, such that $2\delta \tau \approx 0.088$. Using $\tau_{\text{Planck}18} = 0.054$ and $\sigma_{\text{Planck}18} = 0.0073$, this would require

$$\tau_{\rm true} \ge 0.098 = \tau_{\rm Planck18} + 6.0\sigma_{\rm Planck18}$$
 . (2.14)



Figure 4: The historical measurement of the optical depth, τ , from WMAP data [40–44] and Planck [17, 45–48] by year of publication. The horizontal solid (dashed) blue line indicates the central value of τ that would be to move the peak of the DESI+CMB the $\sum m_{\nu}$ posterior from -160 meV (0 meV) to 58 meV.

Similarly, if we take $\tau = 0.051 \pm 0.006$ or $\tau = 0.058 \pm 0.006$ from [47] and [48], we would require shifts of 7.8 σ or 6.7 σ respectively. For comparison, to shift $\sum m_{\nu} = 0$ to 58 meV only requires A_s to be 2.5% larger, which can be accomplished by a $\tau = 0.066$ which is a 1.7 σ upward shift. Both lines are shown in Figure 4 and are consistent with some historical measurements; thus a systematic offset in the more recently inferred values of the optical depth is a plausible explanation for preference for negative neutrino mass. Yet, due to the magnitude of the difference it is unlikely to be the result of a statistical fluctuation.

One of the key challenges with the optical depth is that it is very difficult to measure with ground-based surveys (although it is currently being pursued, for example, by the Cosmology Large Angular Scale Surveyor (CLASS) collaboration [49, 50]). The results of DESI alone point to the need for a confirmation of the Planck measurement of the optical depth, and in principle an improvement to the cosmic variance limit of $\sigma(\tau) = 0.002$. This would be possible with another satellite, such as LiteBird [51]. However, there is the more immediate potential of balloon-based observations which could reach similar levels of sensitivity [52]. Other longer term possibilities include using measurements of crosscorrelations between the CMB and galaxy surveys to eliminate the need for an optical depth measurement [53, 54] or to use measurement the patchy kinetic Sunyaev-Zeldovich effect to constrain the physical model of reionization [55–57], both of which might be possible with CMB-S4 [58].



Figure 5: The historical measurement of the matter density, $\Omega_m h^2$, from WMAP data [40–43] and Planck [17, 45] by year of publication. The black Planck+DESI point is the result of our reanalysis of $\Lambda \text{CDM} + \sum m_{\nu}$ using the same priors as [10]. The horizontal solid (dashed) blue line indicates the central value of $\Omega_m h^2$ that would be required to move the peak of the CMB+DESI $\sum m_{\nu}$ posterior from -160 meV (0 meV) to 58 meV.

Matter Content

The measurement of the matter density ω_m is equally important to the measurement of $\sum m_{\nu}$ as the optical depth. The primary CMB directly determines ω_m through its influence on the height and locations of the acoustic peaks. This is, in part, why the CMB alone is capable of producing very stringent bounds on $\sum m_{\nu}$, e.g. $\sum m_{\nu} < 240 \text{ meV} (95\%)$ from Planck TTTEEE + lensing [17].

Improvements in the measurement to ω_m beyond the CMB has been driven by BAO measurements, most recently with DESI. As shown in Figure 5, the BAO has played a significant role in reducing uncertainty, but has been consistent with the measurements from the CMB data on which the BAO is calibrated. Like the measurement of the optical depth, there was a significant improvement from WMAP to Planck. However, unlike τ , the Planck measurements of ω_m have been stable with the inclusion of more data, including from polarization and the BAO.

The measurement of ω_m needs to be accurate to less than 0.8% in order to permit a reliable measurement of $\sum m_{\nu}$. While this is a high standard, we have the benefit that ω_m will be measured using a number of different CMB surveys that can be combined with several large-scale structure (LSS) surveys. Any large shifts in ω_m due to systematic effects should be different for different surveys and thus from planned measurements alone, we should be able to determine a robust value of ω_m and/or identify systematic issues. This is in sharp contrast to the optical depth, of which Planck is currently the only measurement at the needed accuracy, and it is unclear if near term observations will reproduce or exceed their sensitivity.

It is well known that introducing dynamical dark energy, e.g. in the form of $w_0 \neq -1$ and $w_a \neq 0$, significantly weakens ¹ the neutrino mass constraints [61]. This is for the simple reason that if we allow for more free parameters in the expression for H(z) at low redshifts, we cannot measure ω_m at the accuracy needed to determine $\sum m_{\nu}$. However, this will typically require fairly significant changes to the content and history of the universe. Leaving the content of the universe fixed, we will see that the neutrino mass signal can be explained with changes to the micro-physics in the neutrino and/or dark sector that otherwise leave the rest of cosmological history intact.

CMB Lensing

Weak gravitational lensing of the CMB perturbs the path of photons, so that the apparently location on the sky is perturbed from the true direction $\hat{n}' = \hat{n} + \vec{\alpha}(\hat{n})$, where $\vec{\alpha}(\hat{n})$ is deflection angle [28]. Since the gravitational lensing is time-independent on the scales of observations, the maps of the CMB temperature anistropies (for example) are also modified by the same effect,

$$T_{\text{lensed}}\left(\hat{n}\right) = T_{\text{unlensed}}\left(\hat{n} + \vec{\alpha}(\hat{n})\right) \,. \tag{2.15}$$

The deflection angle is related to the gravitational potential via the lensing potential $\phi(\hat{n})$, via $\vec{\alpha} = \nabla_{\hat{n}} \phi$ and

$$\phi(\hat{n}) \equiv -2 \int_{\eta_{\star}}^{\eta_{0}} \mathrm{d}\eta \frac{\eta - \eta_{\star}}{(\eta_{0} - \eta_{\star})(\eta_{0} - \eta)} \Psi\left((\eta_{0} - \eta)\hat{\mathbf{n}}, \eta\right) , \qquad (2.16)$$

where Ψ is the Weyl potential and η_{\star} is the conformal time of CMB last scattering.

We can understand the main influence of lensing on the CMB by Taylor expanding

$$T_{\text{lensed}}(\hat{n}) \approx T_{\text{unlensed}}(\hat{n}) + \nabla_{\hat{n}}T \cdot \nabla_{\hat{n}}\phi + \mathcal{O}(\phi^2).$$
 (2.17)

For a small patch of sky, we can Fourier transform $\hat{n} \to \vec{\ell}$ so that the dot product is replaced with a convolution

$$T_{\text{lensed}}\left(\vec{\ell}\right) \approx T_{\text{unlensed}}\left(\vec{\ell}\right) - \int \frac{d^2\vec{L}}{2\pi} \vec{L} \cdot (\vec{\ell} - \vec{L}) T_{\text{unlensed}}\left(\vec{\ell} - \vec{L}\right) \phi(\vec{L}) + \mathcal{O}(\phi^2) \,. \tag{2.18}$$

¹Imposing $\sum m_{\nu} > 0$, it has been observed that constraints from current data on neutrino mass can tighten when marginalizing over some models of non-phantom dynamical dark energy [59,60]. While the Cramer-Rao bound requires that the statistical uncertainty must increase, a shift of the central value to more negative values could explain this behavior.

This will induce a non-vanishing correlation between different Fourier modes,

$$\left\langle T_{\text{lensed}} \left(\vec{\ell} \right) T_{\text{lensed}} \left(\vec{L} - \vec{\ell} \right) \right\rangle_{T} = \delta(\vec{L}) C_{\ell}^{TT, \text{unlensed}} + \frac{1}{2\pi} \left[\left(\vec{L} - \vec{\ell} \right) \cdot \vec{L} C_{|\vec{L} - \ell|}^{TT, \text{unlensed}} + \vec{\ell} \cdot \vec{L} C_{\ell}^{TT, \text{unlensed}} \right] \phi(\vec{L}) + \mathcal{O}(\phi^{2}) , \qquad (2.19)$$

where $C_{\ell}^{TT,\text{unlensed}}$ is in the unlensed temperature power spectrum, and the subscript T on the left-hand side refers to an ensemble average over the unlensed CMB temperature realization. As the $\vec{L} \neq 0$ correlations would vanish without lensing, we can reconstruct $\phi(\vec{L})$ from the presence of these correlations [62]. Estimating the CMB lensing power spectrum can therefore be achieved by measuring the temperature four-point function. Lensing also induces a measurable smoothing effect on the acoustic peaks of the CMB power spectrum, from convolving the unlensed power spectrum with the lensing power spectrum at second order.

Once the lensing potential is reconstructed, it can be used to calculate the power spectrum of lensing, remove lensing from the CMB maps [63–65], and/or cross-correlate with other data. For the neutrino mass, the only piece of information we need it the power spectrum of the lensing map $C_L^{\phi\phi}$. This is the same information that is contained in the connected trispectrum of the temperature, as $\phi(\vec{L})$ was determined from a temperature two-point function. As shown in Figure 2, $\sum m_{\nu} = 58$ meV causes a roughly 2-3% suppression of the lensing power, while $\sum m_{\nu} = -160$ meV is a 6-9% enhancement.

The reconstruction of the lensing map is a non-trivial process that could be influenced by other effects that correlate modes in the temperature maps. For example, it is known that the non-Gaussian statistics of unresolved foregrounds can induce biases in these maps [66]. Furthermore, these same correlations are relevant to the covariance of the primary CMB and thus are important for measurements of any other cosmological parameters. Yet, it is also noteworthy that the neutrino mass measurement not sensitive to non-linear effects in the matter power spectrum. Using current CMB data, the lensing map is too noisy to resolve modes that are strongly influenced by non-linear evolution. Yet, even with future data, such as from CMB-S4, these modes can be removed from the analysis with no loss of sensitivity to $\sum m_{\nu}$ [26].

3 Vanishing Neutrino Mass

In this section, we will explore mechanisms for eliminating the signal of $\sum m_{\nu} \geq 58$ meV, while being consistent with $\sum m_{\nu} \geq 0$. The common element of all these models is that we will reduce or eliminate the suppression of power by directly altering the behavior of the neutrinos. In the next section, we will consider changes to the growth of structure beyond just the neutrinos, which could allow for an apparent enhancement of structure, which might be interpreted as $\sum m_{\nu} < 0$.

3.1 Decays

Perhaps the most obvious way to reconcile a cosmological indication of $\sum m_{\nu} = 0$ with the nonzero masses implied by neutrino oscillations is if massive neutrinos decay into massless degrees of freedom on cosmological timescales. While the two heaviest neutrino mass eigenstates are already unstable within the Standard Model, their lifetimes are far greater than the age of the universe $(\tau_{\nu} \propto \langle h \rangle^4 / m_{\nu}^5)$, where $\langle h \rangle \simeq 246$ GeV is the vacuum expectation value of the Higgs field). Neutrino decays on cosmologically relevant timescales would therefore be unambiguous evidence of new physics, above and beyond the origin of neutrino masses.

While decays involving photons are strongly constrained by CMB spectral distortions [67], decays into dark radiation (and either an active or sterile neutrino) are consistent with current limits over a wide range of lifetimes. A lower bound comes from the requirement that the decays and inverse decays of relativistic neutrinos do not prevent free streaming, $\tau_{\nu} \gtrsim 4 \times 10^6 \,\mathrm{s} \,(m_{\nu}/0.05 \,\mathrm{eV})^5$ [68]. On the upper end, the maximum neutrino lifetime that can erase the cosmological signal of neutrino masses depends on the mass spectrum [69–73]. For the minimum masses implied by neutrino oscillations, the lifetime of the massive neutrinos should be roughly an order of magnitude shorter than the age of the universe, $\tau_{\nu} \lesssim 4 \times 10^{16} \,\mathrm{s}$. For the sum of neutrino masses to be observable at KATRIN (sensitive to m_{ν_e} as small as 0.2 eV [74], which translates to $\sum m_{\nu} \sim 0.6 \,\mathrm{eV}$), the maximum lifetime of all the active neutrinos should be around two orders of magnitude smaller, $\tau_{\nu} \lesssim 4 \times 10^{14} \,\mathrm{s}$.

There are a variety of possible decay modes. Two-body decays of massive neutrinos necessarily proceed into a fermion and a boson, with the former either an active or sterile neutrino, and the latter a scalar ϕ or vector Z'. As the masses of the bosons increase, the two-body decay channels close and the bosons instead mediate three-body decays into active and sterile neutrinos. As the viable parameter space for three-body decays is considerably more constrained, here we will restrict our attention to the two-body decays.

In the neutrino mass basis, decays into a (pseudo)scalar arise via couplings of the form

$$\mathcal{L}_{\phi} \supset \frac{\lambda_{ij}}{2} \bar{\nu}_i \nu_j \phi + \frac{\tilde{\lambda}_{ij}}{2} \bar{\nu}_i \gamma_5 \nu_j \phi + \text{h.c.} \qquad (i, j = 1, \dots 4) , \qquad (3.1)$$

where i = 1, 2, 3(4) denote the primarily active (sterile) neutrino mass eigenstates; for definiteness we assume the neutrinos are Majorana. Assuming the lightest active or sterile neutrinos are much lighter than the heavy neutrinos, the corresponding lifetime for decay via the pseudoscalar coupling is [71]

$$\tau(\nu_i \to \nu_j \phi) \simeq 7 \times 10^{17} \,\mathrm{s} \times \left(\frac{0.05 \,\mathrm{eV}}{m_{\nu_i}}\right) \left(\frac{10^{-15}}{\tilde{\lambda}_{ij}^2}\right)^2 \,. \tag{3.2}$$

For two-body decays into active neutrinos to reconcile oscillation splittings with a cosmological measurement of $\sum m_{\nu} = 0$, necessarily $m_{\nu_3} \approx 0.05$ eV. Erasing the energy density in massive neutrinos without spoiling free streaming then implies $4 \times 10^{-15} \leq \lambda, \tilde{\lambda} \leq$ 4×10^{-10} [75–79]. The situation is analogous for decays into sterile neutrinos, although in this case the overall mass scale of active neutrinos may be significantly increased [72].

While the dimensionless couplings required to erase the cosmological neutrino mass signal are small, they are nicely compatible with expectations from UV-complete models. For example, models with spontaneously broken global horizontal lepton flavor symmetries [80] give rise to a goldstone mode coupling to neutrinos as in Eq. (3.1). In such models the off-diagonal pseudoscalar couplings $\tilde{\lambda}_{ij}$ are generated via mixing between heavy sterile and light active neutrinos of order $\tilde{\lambda}_{ij} \simeq \sqrt{m_{\nu_i} m_{\nu_j}}/f$, where f is the scale of spontaneous symmetry breaking. The desired size of $\tilde{\lambda}$ corresponds to 50 MeV $\lesssim f \lesssim 5$ TeV, implying new physics associated with neutrino mass generation around the TeV scale.

Alternately, decays into a vector arise via couplings of the form

$$\mathcal{L}_{Z'} \supset \frac{g_{ij}^L}{2} Z'_{\mu} \bar{\nu}_i \gamma^{\mu} P_L \nu_j + \frac{g_{44}^R}{2} Z'_{\mu} \bar{\nu}_4 \gamma^{\mu} P_R \nu_4 + \text{h.c.} \qquad (i, j = 1, \dots, 4) , \qquad (3.3)$$

which set a lifetime via two-body decays of order

$$\tau(\nu_i \to \nu_j Z') \simeq 7 \times 10^{17} \,\mathrm{s} \times \left(\frac{0.05 \,\mathrm{eV}}{m_{\nu_i}}\right)^3 \left(\frac{m_{Z'}/g_{ij}^L}{50 \,\mathrm{TeV}}\right)^2 \,. \tag{3.4}$$

For two-body decays into active neutrinos to erase the cosmological neutrino mass signal without spoiling free streaming requires $100 \text{ MeV} \leq m_{Z'}/g^L \leq 10 \text{ TeV}$, along with $m_{Z'} \ll m_{\nu_i}$. The situation is analogous for decays into sterile neutrinos, modulo the greater freedom in the active neutrino masses.

As in the scalar case, the dimensionless couplings required to erase the cosmological neutrino mass signal are nicely compatible with expectations from UV-complete models. For instance, in a model with a gauged lepton flavor symmetry such as $U(1)_{L_{\mu}-L_{\tau}}$ broken at a scale f, we have $f = m_{Z'}/g^L$ and the preferred range of decay couplings once again suggests new physics around the TeV scale. The preferred range of couplings and masses is also compatible with current limits, with the most stringent direct bounds $m_{Z'}/g^L >$ 1.3 GeV coming from monolepton + missing energy searches at the LHC [81].

3.2 Annihilation

The cosmological neutrino mass signal may alternately be erased if the cosmological population of massive neutrinos annihilates away into light states at late times [82]. For simplicity, consider the case of a single light (pseudo)scalar coupling to neutrinos, as in Eq. (3.1). Whereas neutrino decays require off-diagonal couplings in the mass basis, annihilation is efficient even when the largest couplings are diagonal. For annihilations to effectively deplete the relic neutrino abundance, the couplings λ , $\tilde{\lambda}$ should be large enough to keep ϕ in thermal equilibrium with neutrinos until after the neutrinos become non-relativistic, at which point the neutrinos annihilate efficiently via $\nu\nu \rightarrow \phi\phi$. The relic neutrino population is effectively erased provided λ , $\tilde{\lambda} \gtrsim 10^{-5}$. However, such large couplings bring ϕ into thermal equilibrium before big bang nucleosynthesis (BBN), and the model is ruled out by a combination of free-streaming requirements and CMB bounds on N_{eff} .

However, mild variations on this scenario remain consistent with current cosmological bounds [83]. One natural possibility is for the active neutrinos to coannihilate into sterile neutrinos via a scalar or pseudoscalar ϕ through the couplings in Eq. (3.1). Avoiding efficient coannihilation while neutrinos are still in thermal equilibrium implies $m_{\phi} \lesssim \text{MeV}$, while ending coannihilation before recombination implies $m_{\phi} \gtrsim \text{eV}$. Within this mass range, efficient conversion requires $\lambda, \tilde{\lambda} \gtrsim 5 \times 10^{-11} \times \left(\frac{m_{\phi}}{\text{keV}}\right)^{1/2}$, while preserving free streaming at recombination requires $\lambda, \tilde{\lambda} \lesssim 5 \times 10^{-3} \times \left(\frac{m_{\phi}}{\text{keV}}\right)$.

The above bounds are based on the direct coupling of active neutrinos to ϕ . These are significantly weakened if the active neutrino couples to ϕ via light right handed neutrinos. In this case, in the early universe, the mixing of relativistic active neutrinos to the right handed neutrino is suppressed by the small neutrino mass, suppressing annihilation at early times. At low redshift, the mixing of non-relativistic neutrinos is unsuppressed, leading to enhanced annihilation that can also explain this signal. We briefly comment on this possibility in Section 3.4.

3.3 Cooling and Heating

The origin of the neutrino mass signal in the matter power spectrum is that the neutrinos are cold enough to redshift like matter, but not cold enough to cluster like matter. Naturally, we could eliminate this signal by either heating or cooling the neutrinos. However, any large change to the temperature would have to come after recombination, as the measurement $N_{\rm eff} = 2.99 \pm 0.33 \ (95\%) \ [17]$ is in precise agreement with the neutrino density predicted by the Standard Model [84–89] and inferred from BBN [90].

Cooling the neutrinos can be an effective strategy if they can be cooled enough to reduce the free-streaming scale below the nonlinear scale, or equivalently $k_{\rm fs} > k_{\rm NL} = \mathcal{O}(1) h \,\mathrm{Mpc}^{-1}$. Recall that the free-streaming scale is defined by

$$k_{\rm fs} = \sqrt{\frac{3}{2}} \frac{aH}{c_{\nu}} , \qquad (3.5)$$

where the neutrino speed in the Standard Model is given by

$$c_{\nu} = \frac{\langle p_{\nu} \rangle}{m_{\nu}} = \frac{3T_{\nu}}{m_{\nu}} \approx 1.0 \times 10^{-2} \left(\frac{50 \,\mathrm{meV}}{m_{\nu}}\right) \,(1+z) \,,$$
 (3.6)

As a result, the free-streaming scale as a function of the neutrino temperature is

$$k_{\rm fs}(z) = 0.04 \, h \, \rm Mpc^{-1} \, \left(\frac{\sum m_{\nu}}{58 \, \rm meV}\right) \times \left(\frac{1.95 \, \rm K}{T_{\nu}(z=0)} \frac{1}{1+z}\right) \,. \tag{3.7}$$

The role of the z-dependence puts a somewhat non-trivial requirement on T_{ν} . At a minimum, if we have $k_{\rm fs}(z \approx 100) > 0.1 \, h \, {\rm Mpc}^{-1}$, then we could expect the neutrinos to cluster on the scales in the linear regime of our late time observations. Less conservatively, we require $k_{\rm fs}(z=0) > 0.1 \, h \, {\rm Mpc}^{-1}$. Together, these imply we need to cool the neutrinos by a factor of 10 to 1000 at redshifts z < 1000 to avoid the neutrino mass signal.

Solving for the coupled linear evolution of the dark matter, baryon, and neutrinos numerically (see Appendix A), Figure 6 shows the suppression as a function of the neutrino temperature as z = 0, T_{ν} for the minimum sum of neutrino masses, $\sum m_{\nu} = 58$ meV. From these numerical results, we can conclude that $T_{\nu} < \mathcal{O}(1) \times 10^{-2}$ K at z = 0 is sufficient to move the free-streaming signal to the non-linear regime, assuming that neutrino cooling occurs near z = 100.

A natural mechanism for cooling the neutrinos is through interactions with dark matter. The dark matter is cold and therefore is a natural heat sink for the neutrinos. It is straightforward [91] to couple a right-handed neutrino, N, to dark matter, χ at lowredshifts through a light mediator ϕ ,

$$\mathcal{L} \supset g_N \phi NN + g_\chi \phi \chi \chi + m^2 \phi^2 + m_N NN + \lambda h LN + m_\chi \chi \chi .$$
(3.8)

Scattering between the dark matter and neutrinos scales as T_{ν}^{-6} and thus avoids the constraints at earlier times (and higher temperatures) from BBN and the CMB [92].

In order to cool the neutrinos and reproduce the clustering in a $\sum m_{\nu} = 0$ universe, it is important that the scattering between neutrinos and dark matter is ineslatic. This could be achieved through a number of mechanisms such a additional dark radiation coupled to χ or having nearly degenerate states associated with χ (like would occur with atomic dark matter, for example). This allows the dark matter to absorb energy from the neutrinos and allows for T_{ν} to decrease. In the above model, $g_N \sim g_{\chi} \approx 10^{-7}$ is sufficient to bring these two sectors into equilibrium at $z \leq 100$ [91] and any efficient process for absorbing the neutrino's energy would lead to an effecive $\sum m_{\nu} = 0$ signal.

One could also consider the case where χ is a single particle sub-component of the dark matter with total energy fraction f_{χ} . Without any additional light states, the scattering between ν and χ is purely elastic. In this scenario, the effect of the coupling is create a neutrino-dark matter fluid, much like the photon-baryon fluid that fills the universe before



Figure 6: Suppression of $P_m(k)$ for $\sum m_{\nu} = 58$ meV and various neutrino temperatures at redshift zero, $T_{\nu}(z=0)$. As the suppression is a percent level effect, it will only be observable in the linear regime $k < 0.1 h \,\mathrm{Mpc}^{-1}$. We see that cooling to $T_{\nu} < 0.02$ K, or cooling by factor of 100, is sufficient to eliminate the signal of free-streaming neutrinos.

recombination, with a free-streaming scale:

$$k_{\rm fs} \approx 0.05 \, h \, {\rm Mpc}^{-1} \times \left(\frac{f_{\chi} + f_{\nu}}{f_{\nu}} \right)^{1/2} \left(\frac{\sum m_{\nu}}{58 \, {\rm meV}} \right).$$
 (3.9)

The amplitude of the suppression on scales $k \gg k_{\rm fs}$ is proportional to $f_{\chi} + f_{\nu}$, the total energy fraction in this fluid. As a result, even if we could couple to all the dark matter so that $k_{\rm fs} = 0.8 \, h \, {\rm Mpc}^{-1}$, the suppression is large enough to be constrained by the Lyman- α forest [93–96] or counts of satellite galaxies [97,98].

Heating the neutrinos to avoid the suppression of matter clustering requires that the neutrino speed, shown in Eq. (3.6), remain near unity throughout cosmic history. This could be achieved by increasing T_{ν} by a factor of ~ 100 in the regime $1000 \gtrsim z \gtrsim 100$; however, this would correspond to increasing the energy density of the cosmic neutrino background ($C\nu B$) by at least the same factor (assuming no change to the number density of neutrinos). The extra energy density acquired by neutrinos needs to be transferred from another component, with the dark matter serving as the natural candidate during the matter-dominated era. A transfer of energy from the dark matter to the $C\nu B$ will have similar cosmological effects as models of dark matter decaying into dark radiation, which are subject to constraints from observations of the matter power spectrum and of the CMB that arise from a larger late-time integrated Sachs-Wolfe effect as compared to a standard cosmological history [99–102]. Current constraints set an upper limit of about 4% of dark matter decaying into radiation after recombination [102], comparable to the fraction of energy density that would need to be transferred from dark matter to heat the $C\nu B$ in order to keep neutrinos relativistic until the present time.

3.4 Time Varying Mass

The tension between the DESI data and the laboratory measurement of neutrino masses can also be alleviated if the mass of the neutrino is not a constant in either time or space. For example, it might be the case that neutrinos had a smaller mass in the early universe (until around $z \sim 10$) but then subsequently had their mass change by $\mathcal{O}(1)$, as suggested in [103–105]. Alternately, it could be the case that the neutrino is a chameleon which acquires a larger mass near high density matter [106] but is otherwise lighter in the low density of the cosmos that is relevant to DESI and CMB lensing. For the purposes of illustration of this concept, in this paper, we study the possibility that the neutrino mass evolved in time and leave further exploration of potential chameleonic nature of neutrinos for future work.

To realize the phenomenology of lower mass neutrinos that become more massive around $z \sim 10$, consider the following terms of the Lagrangian (3.8):

$$\mathcal{L} \supset yhLN + g_N \phi NN + m^2 \phi^2 . \tag{3.10}$$

We take the Yukawa coupling $y \sim \frac{10 \text{ meV}}{\langle h \rangle}$ so that the neutrino's Dirac mass is comparable to the current neutrino mass ~ 10 meV (per neutrino). Observe that when $g_N \phi \gg y \langle h \rangle$ the phenomenology is identical to that of the conventional "see-saw" mechanism and thus the neutrino mass will be light. When $g_N \phi \ll y \langle h \rangle$, the Dirac mass will dominate and equal the desired present day value. The cosmological evolution of $\langle \phi \rangle$ naturally leads to such a change due to the fact that $\langle \phi \rangle$ is sourced by the C ν B, whose number density drops as the universe expands.

To illustrate this dynamic, let us pick some example numbers. Suppose we assume that the neutrino mass was around ~ 1 meV in the early universe. Such neutrinos would be relativistic until $z \sim 10$. When they are relativistic, the C ν B sources $\langle \phi \rangle \sim g_N \frac{\text{meV}^3}{m^2}$ [91], independent of the temperature of the neutrinos. Once the neutrinos become nonrelativistic, $\langle \phi \rangle$ scales with the number density of the C ν B and we get $\langle \phi \rangle \sim g_N \frac{T^3}{m^2}$. When $\langle \phi \rangle$ drops, the neutrinos become more massive, approaching their Dirac mass. The main constraint on this scenario is the bound $g_N \lesssim 5 \times 10^{-8}$ in order to ensure that the neutrinos do not annihilate into ϕ when they are light (in fact, if they do, the situation reduces to the annihilation scenarios considered earlier). Setting $g_N \sim 10^{-8}$ and $m \sim 10^{-12}$ eV, we see that at early times the neutrino mass is around ~ 1 meV. These neutrinos become non-relativistic around $z \sim 10$. The subsequent drop in $\langle \phi \rangle$ raises the neutrino mass to around ~ 20 meV today (per neutrino).

3.5 Mirror Sectors and Relation to the Hubble Tension

The deviation from ΛCDM for $\sum m_{\nu}$ is roughly consistent with the suggestion that new physics might only impact dimensionful parameters [107,108]. The CMB and LSS directly

measure dimensionless quantities (angles, redshifts) and thus are not directly related to dimensionful quantities like H_0 and $\sum m_{\nu}$. This idea was put forward in Refs. [107, 108] to explain the Hubble tension. They realized this concept by introducing a mirror of the Standard Model in the dark sector, such that the gravitational signals remained unchanged but the Standard Model densities could be rescaled.

Naturally, such a model could also easily explain the apparent $\sum m_{\nu} \approx 0$, by having massless neutrinos in the hidden sector. This would leave the other gravitational signals unchanged, but reduce the total gravitational influence of the Standard Model neutrinos. This dilutes $\sum m_{\nu}$ by the fraction of matter in the hidden sector to the mirror sector, and thus requires the Standard Model to be a small component of the total matter density. Unlike some of the other solutions to $\sum m_{\nu}$, this requires an order one change to the universe and thus is difficult to make compatible with all observations. For example, BBN is sensitive to the physical baryon density and thus is not compatible with the simplest implementations of this idea.

Interestingly, the suggestion that there could be multiple copies of the Standard Model with different mass parameters is a natural consequence of several recent mechanisms for solving the hierarchy problem [109–111]. However, these hidden sector typically increase $N_{\rm eff} > 3.044$ and $\sum m_{\nu} > 58$ meV. Without fine tuning these models to take the form of those described in Refs. [107, 108], observations that favor $\sum m_{\nu} < 58$ meV severely constrain these models.

4 Negative "Neutrino Mass"

The possibility of an apparent measurement with $\sum m_{\nu} < 0$ would be most naturally explained by an increase in the amount of clustering in the late universe, or at least an apparent increase as measured through gravitational lensing of the CMB. Even if neutrinos were truly massless $\sum m_{\nu} = 0$, this would require a change to the formation of structure or the statistical properties of the CMB. Such a mechanism could also erase the signal from conventional massive neutrinos and thus need not involve a change to the neutrino sector at all. In this section, we will explore representative examples of how this signal could arise. We will consider physically increasing the amount of clustering through a new long range force, and creating a apparent increase in lensing through changes to the statistics of the primordial density fluctuations. Both classes of ideas lead to observable consequences that may already be testable with existing cosmological data.

4.1 Dark Matter with Long Range Forces

The most direct approach to enhancing the clustering of matter is to increase the strength of the long range force between dark matter particles. Such long range forces are very well constrained for ordinary matter, from tests of the equivalence principle [112]. However, if this new force is limited to the dark matter, it would evade most simple equivalence principle tests. It will nonetheless have observable implications for gravitational dynamics that impact structure on galactic [113–116] and cosmological scales [117, 118]. Interestingly, any such force would also violate the single-field consistency conditions for large-scale structure and thus would leave a measurable non-Gaussian imprint on cosmological correlators [119–121], in addition to the any change to the power spectrum.

Following [118, 121, 122], suppose we introduce a massless field φ that couples only to the dark matter with a r^{-2} force similar to Newtonian gravity. This force will modify the momentum conservation equation for the dark matter,

$$\dot{u}_{\rm cdm} + H u_{\rm cdm} = -\frac{1}{a} \left(\Phi + \alpha \varphi \right) , \qquad \nabla^2 \varphi = \alpha 8 \pi G \bar{\rho}_{\rm cdm} \delta_{\rm cdm} .$$
 (4.1)

The resulting linear growth of the dark matter and baryons at $k \gg k_{\rm fs}$ where $\delta_{\nu} = 0$, is described by

$$\ddot{\delta}_{\rm cdm} + \frac{4}{3t} \dot{\delta}_{\rm cdm} = \frac{2}{3t^2} \left[(1 - f_\nu - f_b)(1 + 2\alpha^2)\delta_{\rm cdm} + f_b \delta_{\rm b} \right] , \qquad (4.2)$$

$$\ddot{\delta}_{\rm b} + \frac{4}{3t}\dot{\delta}_{\rm b} = \frac{2}{3t^2} \left[(1 - f_\nu - f_b)\delta_{\rm cdm} + f_b\delta_{\rm b} \right] \,. \tag{4.3}$$

We will define the new growth term as $(1 - f_{\nu} - f_b)(1 + 2\alpha^2) = 1 + \epsilon$, so that ϵ controls the change to the linear evolution. Taking $\delta_{\text{cdm}} = t^{\gamma}$ and $\delta_b = \xi \delta_{\text{cdm}}$ we find

$$\gamma(\gamma - 1) + \frac{4}{3}\gamma - \frac{2}{3}(1 + \epsilon + \xi f_b) = 0 , \qquad \xi\gamma(\gamma - 1) + \xi\frac{4}{3}\gamma - \frac{2}{3}\left(\frac{1 + \epsilon}{1 + 2\alpha^2} + \xi f_b\right) = 0 .$$
(4.4)

To linear order in ϵ and f_b one finds the growing solution

$$\gamma = \frac{2}{3} + \frac{2}{5}(\epsilon + f_b) , \qquad \xi = 1 - (2\alpha^2) .$$
 (4.5)

In the presense of this new long range force, the power spectrum is therefore modified

$$P^{(\epsilon, \sum m_{\nu})}(k \gg k_{\rm fs}, z) \approx \left(1 - 2f_{\nu} + \frac{6}{5}(\epsilon + f_b)\log\frac{1 + z_{\star}}{1 + z}\right) P^{(\epsilon = 0, \sum m_{\nu} = 0)}(k \gg k_{\rm fs}, z) \ . \ (4.6)$$

Here z_{\star} is the redshift where the long-range force becomes important. In most simple models, z_{\star} is the redshift of horizon entry $k = a(z_{\star})H(z_{\star})$. This would make the above signal scale dependent and thus would not mimic the neutrino signal. As a result, cosmological constraints already exclude $\alpha < 0.01$ [117, 118]. Therefore, it is important that z_{\star} is a *k*-independent constant and that the field φ only becomes important at late times. In this case, if we assume the minimum $\sum m_{\nu}$ so that $f_{\nu} = 4 \times 10^{-3}$, as derived in Equation (2.6), we could explain an apparent $\sum m_{\nu} \approx -160$ meV with $\alpha^2 = 7 \times 10^{-3}$.

A phase transition, or some other time- or temperature-dependent physics could change

the mass of φ so that it became massless at $z_{\star} \approx 100$. This would imply equivalence principle violation for the dark matter at later times. Current constraints [113–116] likely require $\alpha^2 \ll 1$ but have not been explored in detail. In addition, this type of equivalence principle violation leaves a number of cosmological [121] and astrophysical signals [123] that could be observed in near-term surveys and experiments. For example, the change to the evolution of matter also alters the galaxy bispectrum in a way that breaks the single-field consistency conditions. This effect is sufficient to measure $\alpha^2 \gtrsim 10^{-3}$ [121] for a quasi-realistic survey.

4.2 Primordial Trispectrum

The trispectrum (four-point function) of the CMB plays two significant roles in the measurement of neutrino mass. First, gravitational lensing induces a connected four point function, and measuring the trispectrum allows us to reconstruct the lensing power spectrum. Secondly, the trispectrum is also what determines the variance of the primary CMB which sets the uncertainty in all our cosmic parameters [124, 125].

A primordial trispectrum of the appropriate shape could mimic the effect of lensing and thus could lead to an apparent increase in the lensing amplitude. Both lensing and primordial trispectra can be measured using the same class of estimators defined in Ref. [126]. Concretely, we could couple the inflaton to an additional field, $\sigma(\vec{x})$, that modulates the amplitude of the adiabatic fluctuations, $\zeta(\vec{x})$, by a term

$$\zeta(\vec{x}) = \zeta_{\rm G}(\vec{x}) + \sqrt{\tau_{\rm NL}^{\sigma}} \zeta_{\rm G}(\vec{x}) \sigma(\vec{x}) . \qquad (4.7)$$

where $\zeta_{\rm G}(\vec{x})$ and $\sigma(\vec{x})$ are Gaussian random fields. This modulation leads to a connected trispectrum

$$\left\langle \zeta_{\vec{k}_{1}} \zeta_{\vec{k}_{2}} \zeta_{\vec{k}_{3}} \zeta_{\vec{k}_{4}} \right\rangle' = \tau_{\rm NL}^{\sigma} P_{\zeta}(k_{1}) P_{\zeta}(k_{3}) P_{\sigma}(|\vec{k}_{1} + \vec{k}_{2}|) + \text{permutations}$$

$$\equiv \tau_{\rm NL}^{\sigma} T(\vec{k}_{1}, \vec{k}_{2}, \vec{k}_{3}, \vec{k}_{4}) .$$
 (4.8)

This is not equivalent to the lensing signal because it is a three-dimensional correlation between the modes, rather than two dimensional. Bounds on this kind of non-Gaussianity for a scale invariant σ , $P_{\sigma} \approx P_{\zeta}$, have been derived from the CMB and yield $\tau_{\rm NL}^{\rm local} < 1700$ (95%) [127]. However, if the power spectrum of σ were taken to be scale dependent to be degenerate with the lensing potential, $\phi(\vec{L})$, it would be projected out of that analysis. Following Ref. [128] (see also Refs. [129,130]), we can estimate how correlated the proposed trispectrum would be with the local model using the Fisher matrix,

$$F(T_1, T_2) = V \int \frac{d^3 \vec{k_1} d^3 \vec{k_2} d^3 \vec{k_3} d^3 \vec{k_4}}{(2\pi)^{12}} \frac{\left\langle \zeta_{\vec{k_1}} \zeta_{\vec{k_2}} \zeta_{\vec{k_3}} \zeta_{\vec{k_4}} \right\rangle_1' \left\langle \zeta_{\vec{k_1}} \zeta_{\vec{k_2}} \zeta_{\vec{k_3}} \zeta_{\vec{k_4}} \right\rangle_2'}{P_{\zeta}(k_1) P_{\zeta}(k_2) P_{\zeta}(k_3) P_{\zeta}(k_4)} (2\pi)^3 \delta^3 \left(\sum \vec{k_i} \right) ,$$

$$(4.9)$$

where $V \propto k_{\min}^{-3}$ is the survey volume. The ratio of the off-diagonal to diagonal terms defines the correlation coefficient between $\tau_{\rm NL}^{\sigma}$ and $\tau_{\rm NL}^{\rm local}$, $C(\tau_{\rm NL}^{\sigma}, \tau_{\rm NL}^{\rm local})$, which is approximately

$$C(\tau_{\rm NL}^{\sigma}, \tau_{\rm NL}^{\rm local}) \approx \frac{\int d^3k P_{\sigma}(k) P_{\zeta}(k)}{\sqrt{\int d^3k P_{\zeta}(k)^2} \sqrt{\int d^3k P_{\sigma}(k)^2}} \,. \tag{4.10}$$

To match the CMB lensing power, we should choose $P_{\sigma}(k) \propto P_m(k)$ so that it takes a similar form to the lensing signal. We therefore require $P_{\sigma}(k) \to 0$ as $k \to 0$, $P_{\sigma} \propto k^{-3}$ as $k \to \infty$, and have a maximum at some $k = k_{\star}$. We would then expect the correlation to be suppressed by $C(\tau_{\rm NL}^{\sigma}, \tau_{\rm NL}^{\rm local}) \approx (k_{\rm min}/k_{\star})^3 \ll 1$. In this regard, the shape of $P_{\sigma}(k)$ may not have to be finely tuned to contribute to the observed lensing trispectrum without violating other CMB trispectrum constraints. Other trispectrum shapes, like those considered in Refs. [128, 131] are usually scale invariant and peak in equilateral configurations where $k \sim k_{\rm max}$.

Although this signal would be degenerate with lensing in the CMB, it would be introduce non-Gaussianity in the late universe that could be measured through the galaxy power spectrum [132] (via scale-dependent bias [133,134]) or cross-correlations between the CMB and LSS [135]. CMB lensing is currently measured at 40σ [12–14,136] and therefore a trispectrum mimicking a 2.5%–7.5% shift in the lensing amplitude would visible at the 1–3 σ level. Given that the current constraints on primordial non-Gaussianity from related models are at least an order of magnitude weaker than Planck constraints [137], we do not expect² current galaxy survey data to be sensitive to such a trispectrum. However, data from DESI, Euclid [138], and particularly SPHEREx [139] are expected to be up to an order of magnitude more sensitive than Planck to this type of non-Gaussian signature. Concretely, SPHEREx is expected to be sensitive to $\tau_{\rm NL} = 130$ at 2σ [140] which is roughly 10 times the sensitivity of Planck [131].

A second possibility is that additional contributions to the trispectrum could increase the true uncertainty in cosmic parameters. This could increase the probability that value of A_s determined from the primary CMB is simply a statistical outlier. Specifically, a large primordial trispectrum increases the deviation of parameters from their mean values. One model that achieves such behavior is disorder in single field inflation [141]. In these models, random features in the inflationary potential introduce, on average, a trispectrum that is identical the Gaussian noise but with a larger or smaller amplitude. One can achieve a

²We are not aware of published constraints on $\tau_{\rm NL}$ from current galaxy survey data in which we can directly compare Planck.

similar effect on $\sum m_{\nu}$ [142] from super-sample covariance [143], through a large amplitude of local-term non-Gaussianity (e.g. $\tau_{\rm NL}^{\rm local}$). To be consistent with CMB constraints, the effective amplitude $\tau_{\rm NL}$ would have to be scale-dependent to avoid the direct constraints from the CMB trispectrum.

5 Conclusions

The exclusion of the minimum sum of neutrino masses, from either the inverted or normal hierarchy, is a remarkable statement of the power of cosmological data. At these masses, neutrinos form only a fraction of a percent of the total energy density of the universe. The presence of cosmic neutrinos has been robustly established during the era of nucleosynthesis [90] (BBN) and recombination [17] (CMB), through the measurement of $N_{\rm eff}$ and therefore their small but measurable impact on the late universe was to be expected. As we have no simple path to a direct measurement of cosmic neutrinos on earth, cosmological observations provide a novel window into the universe, capable of revealing new secrets.

The recent BAO measurements from DESI enrich this story. Allowing $\sum m_{\nu} < 0$, as an indication of enhanced of clustering, we find data from CMB+DESI constrains $\sum m_{\nu} =$ -160 ± 90 meV (68%), excluding at about 3σ even the minimum neutrino masses consistent with neutrino oscillation experiments. Yet, we showed that this measurement can be naturally explained by new physics in the neutrino and/or dark sectors that is otherwise weakly constrained by other experiments and observations. A measurement consistent with $\sum m_{\nu} = 0$ could be naturally explained by neutrino decays, cooling, or time-dependent neutrino masses, pointing to new physics coupled to neutrinos and potentially dark matter (sectors). Achieving $\sum m_{\nu} < 0$ requires physics beyond the neutrino sector but could be explained by new long range forces for dark matter or changes to the primordial statistics. Each class of models naturally suggests signals that could be present in existing data or testable with near term experiments or observations.

It is important that the measurement of $\sum m_{\nu}$ from the CMB and DESI is incompatible with a wide range of proposals for BSM physics that are also otherwise unconstrained. Light but massive relics [144] are extremely common in models of BSM physics, including many approaches to the hierarchy problem, explanations of dark matter, models including light gravitinos [145, 146], etc. These necessarily contribute positively to N_{eff} and $\sum m_{\nu}$ and thus would further exacerbate the tension with the minimum sum of neutrino masses. As a result, any such model would have to incorporate additional physics, of the kind discussed in this paper, in addition to the new physics relevant to these problems. It is interesting that our results from neutrino decay point to a possible origin from new physics at 10-100 TeV, which could provide a common origin for both effects.

Acknowledgements We are grateful to Kim Berghaus, Tim Cohen, Raphael Flauger, George Fuller, Peter Graham, Jiashu Han, Colin Hill, Mustapha Ishak, Thomas Kon-

standin, Tongyan Lin, and Ben Wallisch for helpful discussions. NC is supported by the US Department of Energy under grant DE-SC0011702. DG is supported by the US Department of Energy (DOE), Office of Science, National Quantum Information Science Research Centers, Superconducting Quantum Materials and Systems Center (SQMS) under Contract No. DE-AC02-07CH11359. S.R. is also supported in part by the U.S. National Science Foundation (NSF) under Grant No. PHY-1818899, the Simons Investigator Grant No. 827042, and by the DOE under a QuantISED grant for MAGIS and Fermilab. JM is supported by the US Department of Energy under grant DE-SC0010129. Computational resources for this research were provided by SMU's Center for Research Computing. We acknowledge the use of CAMB [30], CLASS [32], IPython [147], and the Python packages Matplotlib [148], NumPy [149], and SciPy [150].

A The Suppression of Clustering

In this appendix, we review the calculation of the linear growth of structure in a universe with massive neutrinos. This calculation explains the suppression of small scale power due to neutrino free streaming, which is the dominant cosmological signal responsible for the constraints on $\sum m_{\nu}$.

Following [27], we define the density contrasts of the dark matter and baryons as $\delta_{\rm cb} = \frac{\delta \rho_{\rm cdm} + \delta \rho_{\rm b}}{\bar{\rho}_{\rm cdm} + \bar{\rho}_{\rm b}}$, and the neutrinos, $\delta_{\nu} = \frac{\delta \rho_{\nu}}{\bar{\rho}_{\nu}}$. Energy and momentum conservation of these species after recombination is then described by the coupled equations

$$\dot{\delta}_{\rm cb}(\vec{k},t) - a^{-1}k^2 u_{\rm cb} = 0 , \qquad \dot{\delta}_{\nu}(\vec{k},t) - a^{-1}k^2 u_{\nu} = 0 , \qquad (A.1)$$

and

$$\dot{u}_{\rm cb} + H u_{\rm cb} = -\frac{1}{a} \Phi , \qquad \dot{u}_{\nu} + H u_{\nu} = -\frac{1}{a} \Phi - \frac{c_{\nu}^2}{a} \delta_{\nu} .$$
 (A.2)

Here we have defined the scalar velocity potential u_i for each species as $\vec{v}_i = \vec{\nabla} u_i$. Finally, Φ is the Newtonian gravitational potential, which obeys

$$\nabla^2 \Phi = 4\pi G \left(\bar{\rho}_{cb} \delta_{cb} + \bar{\rho}_{\nu} \delta_{\nu} \right) . \tag{A.3}$$

In a matter dominated universe, $H^2 \propto a^{-3}$ which implies that $a(t) \propto t^{2/3}$ and $\bar{\rho}_{\rm m} \propto t^{-2}$. Differentiating these equations allows us to eliminate the velocity potential to find two second-order equations

$$\ddot{\delta}_{\rm cb} + \frac{4}{3t}\dot{\delta}_{\rm cb} = \frac{2}{3t^2} \left[f_\nu \delta_\nu + (1 - f_\nu)\delta_{\rm cb} \right] , \qquad (A.4)$$

$$\ddot{\delta}_{\nu} + \frac{4}{3t}\dot{\delta}_{\nu} = -\frac{2\alpha}{3t^2}\delta_{\nu} + \frac{2}{3t^2}\left[f_{\nu}\delta_{\nu} + (1 - f_{\nu})\delta_{\rm cb}\right] , \qquad (A.5)$$

where

$$\alpha \equiv \frac{3k^2 c_\nu^2 t^2}{2a^2} = \frac{k^2}{k_{\rm fs}^2} , \quad c_\nu \equiv \frac{\langle p_\nu \rangle}{m_\nu} , \quad f_\nu \equiv \frac{\Omega_\nu}{\Omega_{\rm m}} . \tag{A.6}$$

From here, one can solve these equations numerically to understand the influence of the neutrinos on the matter fluctuations in the linear regime.

In the regime $\alpha \gg 1$, it easy to understand the solutions as follows: the homogeneous equation for δ_{ν} (i.e. $\delta_{\rm cb} \approx 0$) can be solved to find that $\delta_{\nu} \propto t^{-1/6} \to 0$ as $t \to \infty$. We can also solve the inhomogeneous equation with $\delta_{\nu} = \xi \delta_{\rm cb}$ to find $\xi \propto 1/\alpha \to 0$. Therefore we can focus on $\delta_{\rm cb}$ with $\delta_{\nu} = 0$. Taking the ansatz $\delta_{\rm cb} = t^{\gamma}$ and $\delta_{\nu} = 0$, we get

$$\gamma(\gamma - 1) + \frac{4}{3}\gamma - \frac{2(1 - f_{\nu})}{3} = 0 \to \gamma = \frac{2}{3} - \frac{2}{5}f_{\nu} + \mathcal{O}(f_{\nu}^2)$$
(A.7)

where we kept only the growing solution with $\gamma > 0$. In a matter-dominated universe, $H^2 \propto a^{-3}$ which implies that $a(t) \propto t^{2/3}$, and therefore

$$\delta_{\rm cb}(\vec{k},t) \approx \delta_{\rm cb}(\vec{k},t_{\nu})a(t)^{1-3f_{\nu}/5} , \qquad (A.8)$$

where $1 + z_{\nu} = a(t_{\nu})^{-1}$. Finally, since $\bar{\rho}_m = \bar{\rho}_{cb} + \bar{\rho}_{\nu}$, the total matter density contrast $\delta_m = \delta \rho_m / \bar{\rho}_m$ is given by

$$\delta_m(\vec{k}, t) = \frac{\delta\rho_{\rm cb} + \delta\rho_{\nu}}{\bar{\rho}_m} = \delta_{\rm cb}(\vec{k}, t) \approx (1 - f_{\nu})\delta_{\rm cb}(\vec{k}, t_{\nu})a(t)^{1 - 3f_{\nu}/5} \\ \approx \delta_{\rm cb} \left(1 - f_{\nu} - \frac{3}{5}f_{\nu}\log\frac{1 + z_{\nu}}{1 + z}\right) .$$
(A.9)

This gives rise to the suppression of the power spectrum

$$P^{(\sum m_{\nu})}(k \gg k_{\rm fs}, z) \approx \left(1 - 2f_{\nu} - \frac{6}{5}f_{\nu}\log\frac{1+z_{\nu}}{1+z}\right)P^{(\sum m_{\nu}=0)}(k \gg k_{\rm fs}, z) .$$
(A.10)

In this regard, we see that the suppression is a straightforward consequence of the linear evolution.

References

- J. Lesgourgues and S. Pastor, "Massive neutrinos and cosmology," *Phys. Rept.* 429 (2006) 307–379, arXiv:astro-ph/0603494.
- [2] Topical Conveners: K.N. Abazajian, J.E. Carlstrom, A.T. Lee Collaboration, K. N. Abazajian *et al.*, "Neutrino Physics from the Cosmic Microwave Background and Large Scale Structure," *Astropart. Phys.* 63 (2015) 66-80, arXiv:1309.5383 [astro-ph.CO].
- [3] C. Dvorkin *et al.*, "Neutrino Mass from Cosmology: Probing Physics Beyond the Standard Model," arXiv:1903.03689 [astro-ph.CO].
- [4] Particle Data Group Collaboration, P. A. Zyla *et al.*, "Review of Particle Physics," *PTEP* 2020 no. 8, (2020) 083C01.
- [5] A. Font-Ribera, P. McDonald, N. Mostek, B. A. Reid, H.-J. Seo, and A. Slosar, "DESI and other dark energy experiments in the era of neutrino mass measurements," *JCAP* 05 (2014) 023, arXiv:1308.4164 [astro-ph.CO].
- [6] CMB-S4 Collaboration, K. N. Abazajian *et al.*, "CMB-S4 Science Book, First Edition," arXiv:1610.02743 [astro-ph.CO].
- [7] **DESI** Collaboration, A. Aghamousa *et al.*, "The DESI Experiment Part I: Science, Targeting, and Survey Design," arXiv:1611.00036 [astro-ph.IM].
- [8] M. Kaplinghat, L. Knox, and Y.-S. Song, "Determining neutrino mass from the CMB alone," *Phys. Rev. Lett.* 91 (2003) 241301, arXiv:astro-ph/0303344.
- [9] Z. Pan and L. Knox, "Constraints on neutrino mass from Cosmic Microwave Background and Large Scale Structure," Mon. Not. Roy. Astron. Soc. 454 no. 3, (2015) 3200–3206, arXiv:1506.07493 [astro-ph.CO].
- [10] DESI Collaboration, A. G. Adame *et al.*, "DESI 2024 VI: Cosmological Constraints from the Measurements of Baryon Acoustic Oscillations," arXiv:2404.03002 [astro-ph.CO].
- [11] Planck Collaboration, N. Aghanim *et al.*, "Planck 2018 results. V. CMB power spectra and likelihoods," *Astron. Astrophys.* 641 (2020) A5, arXiv:1907.12875 [astro-ph.CO].
- [12] J. Carron, M. Mirmelstein, and A. Lewis, "CMB lensing from Planck PR4 maps," JCAP 09 (2022) 039, arXiv:2206.07773 [astro-ph.CO].

- [13] ACT Collaboration, F. J. Qu et al., "The Atacama Cosmology Telescope: A Measurement of the DR6 CMB Lensing Power Spectrum and Its Implications for Structure Growth," Astrophys. J. 962 no. 2, (2024) 112, arXiv:2304.05202 [astro-ph.CO].
- [14] ACT Collaboration, M. S. Madhavacheril *et al.*, "The Atacama Cosmology Telescope: DR6 Gravitational Lensing Map and Cosmological Parameters," *Astrophys. J.* 962 no. 2, (2024) 113, arXiv:2304.05203 [astro-ph.CO].
- [15] S. Brieden, H. Gil-Marín, and L. Verde, "Model-agnostic interpretation of 10 billion years of cosmic evolution traced by BOSS and eBOSS data," *JCAP* 08 no. 08, (2022) 024, arXiv:2204.11868 [astro-ph.CO].
- [16] N. Palanque-Delabrouille, C. Yèche, N. Schöneberg, J. Lesgourgues, M. Walther, S. Chabanier, and E. Armengaud, "Hints, neutrino bounds and WDM constraints from SDSS DR14 Lyman-α and Planck full-survey data," JCAP 04 (2020) 038, arXiv:1911.09073 [astro-ph.CO].
- [17] Planck Collaboration, N. Aghanim *et al.*, "Planck 2018 results. VI. Cosmological parameters," *Astron. Astrophys.* 641 (2020) A6, arXiv:1807.06209
 [astro-ph.CO]. [Erratum: Astron.Astrophys. 652, C4 (2021)].
- [18] B. Follin, L. Knox, M. Millea, and Z. Pan, "First Detection of the Acoustic Oscillation Phase Shift Expected from the Cosmic Neutrino Background," *Phys. Rev. Lett.* **115** no. 9, (2015) 091301, arXiv:1503.07863 [astro-ph.CO].
- [19] D. Baumann, D. Green, J. Meyers, and B. Wallisch, "Phases of New Physics in the CMB," JCAP 01 (2016) 007, arXiv:1508.06342 [astro-ph.CO].
- [20] D. Baumann, F. Beutler, R. Flauger, D. Green, A. Slosar, M. Vargas-Magaña, B. Wallisch, and C. Yèche, "First constraint on the neutrino-induced phase shift in the spectrum of baryon acoustic oscillations," *Nature Phys.* 15 (2019) 465–469, arXiv:1803.10741 [astro-ph.CO].
- [21] F.-Y. Cyr-Racine and K. Sigurdson, "Limits on Neutrino-Neutrino Scattering in the Early Universe," *Phys. Rev. D* 90 no. 12, (2014) 123533, arXiv:1306.1536 [astro-ph.CO].
- [22] L. Lancaster, F.-Y. Cyr-Racine, L. Knox, and Z. Pan, "A tale of two modes: Neutrino free-streaming in the early universe," JCAP 07 (2017) 033, arXiv:1704.06657 [astro-ph.CO].
- [23] A. He, R. An, M. M. Ivanov, and V. Gluscevic, "Self-Interacting Neutrinos in Light of Large-Scale Structure Data," arXiv:2309.03956 [astro-ph.CO].

- [24] D. Camarena, F.-Y. Cyr-Racine, and J. Houghteling, "Confronting self-interacting neutrinos with the full shape of the galaxy power spectrum," *Phys. Rev. D* 108 no. 10, (2023) 103535, arXiv:2309.03941 [astro-ph.CO].
- [25] D. Camarena and F.-Y. Cyr-Racine, "Absence of concordance in a simple self-interacting neutrino cosmology," arXiv:2403.05496 [astro-ph.CO].
- [26] D. Green and J. Meyers, "Cosmological Implications of a Neutrino Mass Detection," arXiv:2111.01096 [astro-ph.CO].
- [27] D. Green, "Cosmic Signals of Fundamental Physics," PoS TASI2022 (2024) 005, arXiv:2212.08685 [hep-ph].
- [28] A. Lewis and A. Challinor, "Weak gravitational lensing of the CMB," *Phys. Rept.* 429 (2006) 1–65, arXiv:astro-ph/0601594.
- [29] eBOSS Collaboration, S. Alam *et al.*, "Completed SDSS-IV extended Baryon Oscillation Spectroscopic Survey: Cosmological implications from two decades of spectroscopic surveys at the Apache Point Observatory," *Phys. Rev. D* 103 no. 8, (2021) 083533, arXiv:2007.08991 [astro-ph.CO].
- [30] A. Lewis, A. Challinor, and A. Lasenby, "Efficient Computation of CMB Anisotropies in Closed FRW Models," *Astrophys. J.* 538 (2000) 473, arXiv:astro-ph/9911177.
- [31] C. Howlett, A. Lewis, A. Hall, and A. Challinor, "CMB power spectrum parameter degeneracies in the era of precision cosmology," *JCAP* **1204** (2012) 027, arXiv:1201.3654 [astro-ph.CO]. https://arxiv.org/abs/1201.3654.
- [32] D. Blas, J. Lesgourgues, and T. Tram, "The Cosmic Linear Anisotropy Solving System (CLASS) II: Approximation Schemes," JCAP 07 (2011) 034, arXiv:1104.2933 [astro-ph.CO].
- [33] **DESI** Collaboration, A. G. Adame *et al.*, "DESI 2024 IV: Baryon Acoustic Oscillations from the Lyman Alpha Forest," arXiv:2404.03001 [astro-ph.CO].
- [34] **DESI** Collaboration, A. G. Adame *et al.*, "DESI 2024 III: Baryon Acoustic Oscillations from Galaxies and Quasars," arXiv:2404.03000 [astro-ph.CO].
- [35] J. Torrado and A. Lewis, "Cobaya: Code for Bayesian Analysis of hierarchical physical models," JCAP 05 (2021) 057, arXiv:2005.05290 [astro-ph.IM].
- [36] A. Lewis and S. Bridle, "Cosmological parameters from CMB and other data: A Monte Carlo approach," *Phys. Rev.* D66 (2002) 103511, arXiv:astro-ph/0205436 [astro-ph]. https://arxiv.org/abs/astro-ph/0205436.

- [37] A. Lewis, "Efficient sampling of fast and slow cosmological parameters," *Phys. Rev.* D87 no. 10, (2013) 103529, arXiv:1304.4473 [astro-ph.CO]. https://arxiv.org/abs/1304.4473.
- [38] R. M. Neal, "Taking Bigger Metropolis Steps by Dragging Fast Variables," ArXiv Mathematics e-prints (Feb., 2005), math/0502099.
 https://arxiv.org/abs/math/0502099.
- [39] E. Abdalla *et al.*, "Cosmology intertwined: A review of the particle physics, astrophysics, and cosmology associated with the cosmological tensions and anomalies," *JHEAp* 34 (2022) 49–211, arXiv:2203.06142 [astro-ph.CO].
- [40] WMAP Collaboration, C. L. Bennett *et al.*, "First year Wilkinson Microwave Anisotropy Probe (WMAP) observations: Preliminary maps and basic results," *Astrophys. J. Suppl.* 148 (2003) 1–27, arXiv:astro-ph/0302207.
- [41] WMAP Collaboration, D. N. Spergel *et al.*, "Wilkinson Microwave Anisotropy Probe (WMAP) three year results: implications for cosmology," *Astrophys. J. Suppl.* 170 (2007) 377, arXiv:astro-ph/0603449.
- [42] WMAP Collaboration, E. Komatsu *et al.*, "Five-Year Wilkinson Microwave Anisotropy Probe (WMAP) Observations: Cosmological Interpretation," *Astrophys. J. Suppl.* 180 (2009) 330–376, arXiv:0803.0547 [astro-ph].
- [43] WMAP Collaboration, E. Komatsu *et al.*, "Seven-Year Wilkinson Microwave Anisotropy Probe (WMAP) Observations: Cosmological Interpretation," *Astrophys. J. Suppl.* **192** (2011) 18, arXiv:1001.4538 [astro-ph.CO].
- [44] WMAP Collaboration, G. Hinshaw et al., "Nine-Year Wilkinson Microwave Anisotropy Probe (WMAP) Observations: Cosmological Parameter Results," Astrophys. J. Suppl. 208 (2013) 19, arXiv:1212.5226 [astro-ph.CO].
- [45] Planck Collaboration, P. A. R. Ade et al., "Planck 2013 results. XVI. Cosmological parameters," Astron. Astrophys. 571 (2014) A16, arXiv:1303.5076 [astro-ph.CO].
- [46] Planck Collaboration, P. A. R. Ade *et al.*, "Planck 2015 results. XIII. Cosmological parameters," *Astron. Astrophys.* 594 (2016) A13, arXiv:1502.01589 [astro-ph.CO].
- [47] Planck Collaboration, Y. Akrami *et al.*, "*Planck* intermediate results. LVII. Joint Planck LFI and HFI data processing," *Astron. Astrophys.* 643 (2020) A42, arXiv:2007.04997 [astro-ph.CO].

- [48] M. Tristram *et al.*, "Cosmological parameters derived from the final Planck data release (PR4)," *Astron. Astrophys.* 682 (2024) A37, arXiv:2309.10034 [astro-ph.CO].
- [49] T. Essinger-Hileman et al., "CLASS: The Cosmology Large Angular Scale Surveyor," Proc. SPIE Int. Soc. Opt. Eng. 9153 (2014) 91531I, arXiv:1408.4788 [astro-ph.IM].
- [50] J. R. Eimer *et al.*, "CLASS Angular Power Spectra and Map-component Analysis for 40 GHz Observations through 2022," *Astrophys. J.* 963 no. 2, (2024) 92, arXiv:2309.00675 [astro-ph.CO].
- [51] LiteBIRD Collaboration, E. Allys *et al.*, "Probing Cosmic Inflation with the LiteBIRD Cosmic Microwave Background Polarization Survey," *PTEP* 2023 no. 4, (2023) 042F01, arXiv:2202.02773 [astro-ph.IM].
- [52] J. Errard, M. Remazeilles, J. Aumont, J. Delabrouille, D. Green, S. Hanany, B. S. Hensley, and A. Kogut, "Constraints on the Optical Depth to Reionization from Balloon-borne Cosmic Microwave Background Measurements," *Astrophys. J.* 940 no. 1, (2022) 68, arXiv:2206.03389 [astro-ph.CO].
- [53] B. Yu, R. Z. Knight, B. D. Sherwin, S. Ferraro, L. Knox, and M. Schmittfull, "Toward neutrino mass from cosmology without optical depth information," *Phys. Rev. D* 107 no. 12, (2023) 123522, arXiv:1809.02120 [astro-ph.CO].
- [54] T. Brinckmann, D. C. Hooper, M. Archidiacono, J. Lesgourgues, and T. Sprenger, "The promising future of a robust cosmological neutrino mass measurement," *JCAP* 01 (2019) 059, arXiv:1808.05955 [astro-ph.CO].
- [55] K. M. Smith and S. Ferraro, "Detecting Patchy Reionization in the Cosmic Microwave Background," *Phys. Rev. Lett.* **119** no. 2, (2017) 021301, arXiv:1607.01769 [astro-ph.CO].
- [56] S. Ferraro and K. M. Smith, "Characterizing the epoch of reionization with the small-scale CMB: Constraints on the optical depth and duration," *Phys. Rev. D* 98 no. 12, (2018) 123519, arXiv:1803.07036 [astro-ph.CO].
- [57] M. A. Alvarez, S. Ferraro, J. C. Hill, R. Hlovzek, and M. Ikape, "Mitigating the optical depth degeneracy using the kinematic Sunyaev-Zel'dovich effect with CMB-S4," *Phys. Rev. D* 103 no. 6, (2021) 063518, arXiv:2006.06594 [astro-ph.CO].
- [58] K. Abazajian *et al.*, "CMB-S4 Science Case, Reference Design, and Project Plan," arXiv:1907.04473 [astro-ph.IM].

- [59] S. Vagnozzi, S. Dhawan, M. Gerbino, K. Freese, A. Goobar, and O. Mena, "Constraints on the sum of the neutrino masses in dynamical dark energy models with w(z) ≥ -1 are tighter than those obtained in ΛCDM," *Phys. Rev. D* 98 no. 8, (2018) 083501, arXiv:1801.08553 [astro-ph.CO].
- [60] K. V. Berghaus, J. A. Kable, and V. Miranda, "Quantifying Scalar Field Dynamics with DESI 2024 Y1 BAO measurements," arXiv:2404.14341 [astro-ph.CO].
- [61] R. Allison, P. Caucal, E. Calabrese, J. Dunkley, and T. Louis, "Towards a cosmological neutrino mass detection," *Phys. Rev. D* 92 no. 12, (2015) 123535, arXiv:1509.07471 [astro-ph.CO].
- [62] W. Hu and T. Okamoto, "Mass reconstruction with cmb polarization," Astrophys. J. 574 (2002) 566-574, arXiv:astro-ph/0111606.
- [63] U. Seljak and C. M. Hirata, "Gravitational lensing as a contaminant of the gravity wave signal in CMB," *Phys. Rev. D* 69 (2004) 043005, arXiv:astro-ph/0310163.
- [64] D. Green, J. Meyers, and A. van Engelen, "CMB Delensing Beyond the B Modes," JCAP 12 (2017) 005, arXiv:1609.08143 [astro-ph.CO].
- [65] S. C. Hotinli, J. Meyers, C. Trendafilova, D. Green, and A. van Engelen, "The benefits of CMB delensing," *JCAP* 04 no. 04, (2022) 020, arXiv:2111.15036 [astro-ph.CO].
- [66] A. van Engelen, S. Bhattacharya, N. Sehgal, G. P. Holder, O. Zahn, and D. Nagai, "CMB Lensing Power Spectrum Biases from Galaxies and Clusters using High-angular Resolution Temperature Maps," *Astrophys. J.* 786 (2014) 13, arXiv:1310.7023 [astro-ph.CO].
- [67] J. L. Aalberts *et al.*, "Precision constraints on radiative neutrino decay with CMB spectral distortion," *Phys. Rev. D* 98 (2018) 023001, arXiv:1803.00588 [astro-ph.CO].
- [68] G. Barenboim, J. Z. Chen, S. Hannestad, I. M. Oldengott, T. Tram, and Y. Y. Wong, "Invisible neutrino decay in precision cosmology," *JCAP* 03 (2021) 087, arXiv:2011.01502 [astro-ph.CO].
- [69] Z. Chacko, A. Dev, P. Du, V. Poulin, and Y. Tsai, "Cosmological Limits on the Neutrino Mass and Lifetime," JHEP 04 (2020) 020, arXiv:1909.05275 [hep-ph].
- [70] Z. Chacko, A. Dev, P. Du, V. Poulin, and Y. Tsai, "Determining the Neutrino Lifetime from Cosmology," *Phys. Rev. D* 103 no. 4, (2021) 043519, arXiv:2002.08401 [astro-ph.CO].

- [71] M. Escudero, J. Lopez-Pavon, N. Rius, and S. Sandner, "Relaxing Cosmological Neutrino Mass Bounds with Unstable Neutrinos," *JHEP* 12 (2020) 119, arXiv:2007.04994 [hep-ph].
- [72] G. Franco Abellán, Z. Chacko, A. Dev, P. Du, V. Poulin, and Y. Tsai, "Improved cosmological constraints on the neutrino mass and lifetime," *JHEP* 08 (2022) 076, arXiv:2112.13862 [hep-ph].
- [73] M. Escudero, T. Schwetz, and J. Terol-Calvo, "A seesaw model for large neutrino masses in concordance with cosmology," JHEP 02 (2023) 142, arXiv:2211.01729 [hep-ph].
- [74] KATRIN Collaboration, M. Aker et al., "Direct neutrino-mass measurement with sub-electronvolt sensitivity," *Nature Phys.* 18 no. 2, (2022) 160–166, arXiv:2105.08533 [hep-ex].
- [75] Y. Farzan, "Bounds on the coupling of the Majoron to light neutrinos from supernova cooling," *Phys. Rev. D* 67 (2003) 073015, arXiv:hep-ph/0211375.
- [76] Z. Chacko, L. J. Hall, T. Okui, and S. J. Oliver, "CMB signals of neutrino mass generation," *Phys. Rev. D* 70 (2004) 085008, arXiv:hep-ph/0312267.
- [77] A. Friedland, K. M. Zurek, and S. Bashinsky, "Constraining Models of Neutrino Mass and Neutrino Interactions with the Planck Satellite," arXiv:0704.3271 [astro-ph].
- [78] M. Archidiacono and S. Hannestad, "Updated constraints on non-standard neutrino interactions from Planck," JCAP 07 (2014) 046, arXiv:1311.3873 [astro-ph.CO].
- [79] D. Baumann, D. Green, and B. Wallisch, "New Target for Cosmic Axion Searches," *Phys. Rev. Lett.* **117** no. 17, (2016) 171301, arXiv:1604.08614 [astro-ph.CO].
- [80] G. B. Gelmini and J. W. F. Valle, "Fast Invisible Neutrino Decays," *Phys. Lett. B* 142 (1984) 181–187.
- [81] M. Ekhterachian, A. Hook, S. Kumar, and Y. Tsai, "Bounds on gauge bosons coupled to nonconserved currents," *Phys. Rev. D* 104 no. 3, (2021) 035034, arXiv:2103.13396 [hep-ph].
- [82] J. F. Beacom, N. F. Bell, and S. Dodelson, "Neutrinoless universe," *Phys. Rev. Lett.* **93** (2004) 121302, arXiv:astro-ph/0404585.

- [83] Y. Farzan and S. Hannestad, "Neutrinos secretly converting to lighter particles to please both KATRIN and the cosmos," JCAP 02 (2016) 058, arXiv:1510.02201 [hep-ph].
- [84] G. Mangano, G. Miele, S. Pastor, T. Pinto, O. Pisanti, and P. D. Serpico, "Relic neutrino decoupling including flavor oscillations," *Nucl. Phys. B* 729 (2005) 221-234, arXiv:hep-ph/0506164.
- [85] M. Escudero Abenza, "Precision early universe thermodynamics made simple: N_{eff} and neutrino decoupling in the Standard Model and beyond," JCAP 05 (2020) 048, arXiv:2001.04466 [hep-ph].
- [86] K. Akita and M. Yamaguchi, "A precision calculation of relic neutrino decoupling," JCAP 08 (2020) 012, arXiv:2005.07047 [hep-ph].
- [87] J. Froustey, C. Pitrou, and M. C. Volpe, "Neutrino decoupling including flavour oscillations and primordial nucleosynthesis," JCAP 12 (2020) 015, arXiv:2008.01074 [hep-ph].
- [88] J. J. Bennett, G. Buldgen, P. F. De Salas, M. Drewes, S. Gariazzo, S. Pastor, and Y. Y. Y. Wong, "Towards a precision calculation of N_{eff} in the Standard Model II: Neutrino decoupling in the presence of flavour oscillations and finite-temperature QED," JCAP 04 (2021) 073, arXiv:2012.02726 [hep-ph].
- [89] J. R. Bond, G. M. Fuller, E. Grohs, J. Meyers, and M. J. Wilson, "Cosmic Neutrino Decoupling and its Observable Imprints: Insights from Entropic-Dual Transport," arXiv:2403.19038 [astro-ph.CO].
- [90] B. D. Fields, K. A. Olive, T.-H. Yeh, and C. Young, "Big-Bang Nucleosynthesis after Planck," *JCAP* 03 (2020) 010, arXiv:1912.01132 [astro-ph.CO].
 [Erratum: JCAP 11, E02 (2020)].
- [91] D. Green, D. E. Kaplan, and S. Rajendran, "Neutrino interactions in the late universe," JHEP 11 (2021) 162, arXiv:2108.06928 [hep-ph].
- [92] K. M. Nollett and G. Steigman, "BBN And The CMB Constrain Neutrino Coupled Light WIMPs," *Phys. Rev. D* 91 no. 8, (2015) 083505, arXiv:1411.6005 [astro-ph.CO].
- [93] SDSS Collaboration, P. McDonald *et al.*, "The Lyman-alpha forest power spectrum from the Sloan Digital Sky Survey," *Astrophys. J. Suppl.* 163 (2006) 80–109, arXiv:astro-ph/0405013.

- [94] M. Viel, J. Lesgourgues, M. G. Haehnelt, S. Matarrese, and A. Riotto, "Constraining warm dark matter candidates including sterile neutrinos and light gravitinos with WMAP and the Lyman-alpha forest," *Phys. Rev. D* 71 (2005) 063534, arXiv:astro-ph/0501562.
- [95] A. Boyarsky, J. Lesgourgues, O. Ruchayskiy, and M. Viel, "Lyman-alpha constraints on warm and on warm-plus-cold dark matter models," *JCAP* 05 (2009) 012, arXiv:0812.0010 [astro-ph].
- [96] W. L. Xu, C. Dvorkin, and A. Chael, "Probing sub-GeV Dark Matter-Baryon Scattering with Cosmological Observables," *Phys. Rev. D* 97 no. 10, (2018) 103530, arXiv:1802.06788 [astro-ph.CO].
- [97] E. O. Nadler, V. Gluscevic, K. K. Boddy, and R. H. Wechsler, "Constraints on Dark Matter Microphysics from the Milky Way Satellite Population," *Astrophys. J. Lett.* 878 no. 2, (2019) 32, arXiv:1904.10000 [astro-ph.CO]. [Erratum: Astrophys.J.Lett. 897, L46 (2020), Erratum: Astrophys.J. 897, L46 (2020)].
- [98] K. Maamari, V. Gluscevic, K. K. Boddy, E. O. Nadler, and R. H. Wechsler, "Bounds on velocity-dependent dark matter-proton scattering from Milky Way satellite abundance," *Astrophys. J. Lett.* **907** no. 2, (2021) L46, arXiv:2010.02936 [astro-ph.CO].
- [99] L. Kofman, D. Pogosyan, and A. A. Starobinsky, "The large scale microwave backbround anisotropy in unstable cosmologies," Sov. Astron. Lett. 12 (1986) 175–179.
- [100] S. De Lope Amigo, W. M.-Y. Cheung, Z. Huang, and S.-P. Ng, "Cosmological Constraints on Decaying Dark Matter," JCAP 06 (2009) 005, arXiv:0812.4016 [hep-ph].
- [101] B. Audren, J. Lesgourgues, G. Mangano, P. D. Serpico, and T. Tram, "Strongest model-independent bound on the lifetime of Dark Matter," *JCAP* 12 (2014) 028, arXiv:1407.2418 [astro-ph.CO].
- [102] V. Poulin, P. D. Serpico, and J. Lesgourgues, "A fresh look at linear cosmological constraints on a decaying dark matter component," *JCAP* 08 (2016) 036, arXiv:1606.02073 [astro-ph.CO].
- [103] R. Fardon, A. E. Nelson, and N. Weiner, "Dark energy from mass varying neutrinos," JCAP 10 (2004) 005, arXiv:astro-ph/0309800.
- [104] C. S. Lorenz, L. Funcke, E. Calabrese, and S. Hannestad, "Time-varying neutrino mass from a supercooled phase transition: current cosmological constraints and

impact on the Ω_m - σ_8 plane," *Phys. Rev. D* **99** no. 2, (2019) 023501, arXiv:1811.01991 [astro-ph.CO].

- [105] C. S. Lorenz, L. Funcke, M. Löffler, and E. Calabrese, "Reconstruction of the neutrino mass as a function of redshift," *Phys. Rev. D* 104 no. 12, (2021) 123518, arXiv:2102.13618 [astro-ph.CO].
- [106] H. Davoudiasl, G. Mohlabeng, and M. Sullivan, "Galactic Dark Matter Population as the Source of Neutrino Masses," *Phys. Rev. D* 98 no. 2, (2018) 021301, arXiv:1803.00012 [hep-ph].
- [107] F.-Y. Cyr-Racine, F. Ge, and L. Knox, "Symmetry of Cosmological Observables, a Mirror World Dark Sector, and the Hubble Constant," *Phys. Rev. Lett.* **128** no. 20, (2022) 201301, arXiv:2107.13000 [astro-ph.CO].
- [108] F. Ge, F.-Y. Cyr-Racine, and L. Knox, "Scaling transformations and the origins of light relics constraints from cosmic microwave background observations," *Phys. Rev. D* 107 no. 2, (2023) 023517, arXiv:2210.16335 [astro-ph.CO].
- [109] N. Arkani-Hamed, T. Cohen, R. T. D'Agnolo, A. Hook, H. D. Kim, and D. Pinner, "Solving the Hierarchy Problem at Reheating with a Large Number of Degrees of Freedom," *Phys. Rev. Lett.* **117** no. 25, (2016) 251801, arXiv:1607.06821 [hep-ph].
- [110] Z. Chacko, N. Craig, P. J. Fox, and R. Harnik, "Cosmology in Mirror Twin Higgs and Neutrino Masses," JHEP 07 (2017) 023, arXiv:1611.07975 [hep-ph].
- [111] Z. Chacko, D. Curtin, M. Geller, and Y. Tsai, "Cosmological Signatures of a Mirror Twin Higgs," JHEP 09 (2018) 163, arXiv:1803.03263 [hep-ph].
- [112] C. M. Will, "The Confrontation between General Relativity and Experiment," Living Rev. Rel. 17 (2014) 4, arXiv:1403.7377 [gr-qc].
- [113] M. Kesden and M. Kamionkowski, "Galilean Equivalence for Galactic Dark Matter," Phys. Rev. Lett. 97 (2006) 131303, arXiv:astro-ph/0606566.
- [114] M. Kesden and M. Kamionkowski, "Tidal Tails Test the Equivalence Principle in the Dark Sector," *Phys. Rev. D* 74 (2006) 083007, arXiv:astro-ph/0608095.
- [115] J. A. Keselman, A. Nusser, and P. J. E. Peebles, "Cosmology with Equivalence Principle Breaking in the Dark Sector," *Phys. Rev. D* 81 (2010) 063521, arXiv:0912.4177 [astro-ph.CO].
- [116] Z. Bogorad, P. W. Graham, and H. Ramani, "Coherent Self-Interactions of Dark Matter in the Bullet Cluster," arXiv:2311.07648 [hep-ph].

- [117] M. Archidiacono, E. Castorina, D. Redigolo, and E. Salvioni, "Unveiling dark fifth forces with linear cosmology," *JCAP* 10 (2022) 074, arXiv:2204.08484 [astro-ph.CO].
- [118] S. Bottaro, E. Castorina, M. Costa, D. Redigolo, and E. Salvioni, "Unveiling dark forces with the Large Scale Structure of the Universe," arXiv:2309.11496 [astro-ph.CO].
- [119] P. Creminelli, J. Noreña, M. Simonović, and F. Vernizzi, "Single-Field Consistency Relations of Large Scale Structure," JCAP 12 (2013) 025, arXiv:1309.3557 [astro-ph.CO].
- [120] P. Creminelli, J. Gleyzes, M. Simonović, and F. Vernizzi, "Single-Field Consistency Relations of Large Scale Structure. Part II: Resummation and Redshift Space," *JCAP* 02 (2014) 051, arXiv:1311.0290 [astro-ph.CO].
- [121] P. Creminelli, J. Gleyzes, L. Hui, M. Simonović, and F. Vernizzi, "Single-Field Consistency Relations of Large Scale Structure. Part III: Test of the Equivalence Principle," JCAP 06 (2014) 009, arXiv:1312.6074 [astro-ph.CO].
- [122] F. Saracco, M. Pietroni, N. Tetradis, V. Pettorino, and G. Robbers, "Non-linear Matter Spectra in Coupled Quintessence," *Phys. Rev. D* 82 (2010) 023528, arXiv:0911.5396 [astro-ph.CO].
- [123] Y. Bai, S. Lu, and N. Orlofsky, "Gravitational Waves From More Attractive Dark Binaries," arXiv:2312.13378 [astro-ph.CO].
- [124] W. Hu, "Angular trispectrum of the CMB," Phys. Rev. D 64 (2001) 083005, arXiv:astro-ph/0105117.
- [125] T. Okamoto and W. Hu, "The angular trispectra of CMB temperature and polarization," *Phys. Rev. D* 66 (2002) 063008, arXiv:astro-ph/0206155.
- [126] D. Hanson and A. Lewis, "Estimators for CMB Statistical Anisotropy," Phys. Rev. D 80 (2009) 063004, arXiv:0908.0963 [astro-ph.CO].
- [127] K. Marzouk, A. Lewis, and J. Carron, "Constraints on τ_{NL} from Planck temperature and polarization," *JCAP* **08** no. 08, (2022) 015, arXiv:2205.14408 [astro-ph.CO].
- [128] K. M. Smith, L. Senatore, and M. Zaldarriaga, "Optimal analysis of the CMB trispectrum," arXiv:1502.00635 [astro-ph.CO].

- [129] K. M. Smith, M. LoVerde, and M. Zaldarriaga, "A universal bound on N-point correlations from inflation," *Phys. Rev. Lett.* **107** (2011) 191301, arXiv:1108.1805 [astro-ph.CO].
- [130] D. Green, Y. Huang, C.-H. Shen, and D. Baumann, "Positivity from Cosmological Correlators," JHEP 04 (2024) 034, arXiv:2310.02490 [hep-th].
- [131] Planck Collaboration, Y. Akrami *et al.*, "Planck 2018 results. IX. Constraints on primordial non-Gaussianity," *Astron. Astrophys.* 641 (2020) A9, arXiv:1905.05697 [astro-ph.CO].
- [132] S. Ferraro and K. M. Smith, "Using large scale structure to measure f_{NL}, g_{NL} and τ_{NL} ," *Phys. Rev. D* **91** no. 4, (2015) 043506, arXiv:1408.3126 [astro-ph.CO].
- [133] J.-O. Gong and S. Yokoyama, "Scale dependent bias from primordial non-Gaussianity with trispectrum," Mon. Not. Roy. Astron. Soc. 417 (2011) 79, arXiv:1106.4404 [astro-ph.CO].
- [134] D. Baumann, S. Ferraro, D. Green, and K. M. Smith, "Stochastic Bias from Non-Gaussian Initial Conditions," JCAP 05 (2013) 001, arXiv:1209.2173 [astro-ph.CO].
- [135] N. Anil Kumar, G. Sato-Polito, M. Kamionkowski, and S. C. Hotinli, "Primordial trispectrum from kinetic Sunyaev-Zel'dovich tomography," *Phys. Rev. D* 106 no. 6, (2022) 063533, arXiv:2205.03423 [astro-ph.CO].
- [136] Planck Collaboration, N. Aghanim et al., "Planck 2018 results. VIII. Gravitational lensing," Astron. Astrophys. 641 (2020) A8, arXiv:1807.06210 [astro-ph.CO].
- [137] D. Green, Y. Guo, J. Han, and B. Wallisch, "Light Fields during Inflation from BOSS and Future Galaxy Surveys," arXiv:2311.04882 [astro-ph.CO].
- [138] EUCLID Collaboration, R. Laureijs et al., "Euclid Definition Study Report," arXiv:1110.3193 [astro-ph.CO].
- [139] SPHEREx Collaboration, O. Doré *et al.*, "Cosmology with the SPHEREX All-Sky Spectral Survey," arXiv:1412.4872 [astro-ph.CO].
- [140] C. Shiveshwarkar, T. Brinckmann, and M. Loverde, "Constraining general multi-field inflation using the SPHEREx all-sky survey," arXiv:2312.15038 [astro-ph.CO].
- [141] D. Green, "Disorder in the Early Universe," JCAP 03 (2015) 020, arXiv:1409.6698 [hep-th].

- [142] M. Forconi, E. Di Valentino, A. Melchiorri, and S. Pan, "A Possible Impact of Non-Gaussianities on Cosmological Constraints in Neutrino Physics," arXiv:2311.04038 [astro-ph.CO].
- [143] M. Takada and W. Hu, "Power Spectrum Super-Sample Covariance," Phys. Rev. D 87 no. 12, (2013) 123504, arXiv:1302.6994 [astro-ph.CO].
- [144] W. L. Xu, J. B. Muñoz, and C. Dvorkin, "Cosmological constraints on light but massive relics," *Phys. Rev. D* 105 no. 9, (2022) 095029, arXiv:2107.09664 [astro-ph.CO].
- [145] T. Moroi, H. Murayama, and M. Yamaguchi, "Cosmological constraints on the light stable gravitino," *Phys. Lett. B* 303 (1993) 289–294.
- [146] K. Osato, T. Sekiguchi, M. Shirasaki, A. Kamada, and N. Yoshida, "Cosmological Constraint on the Light Gravitino Mass from CMB Lensing and Cosmic Shear," *JCAP* 06 (2016) 004, arXiv:1601.07386 [astro-ph.CO].
- [147] F. Pérez and B. Granger, "IPython: A System for Interactive Scientific Computing," *Comput. Sci. Eng.* 9 (2007) 21.
- [148] J. Hunter, "Matplotlib: A 2D Graphics Environment," Comput. Sci. Eng. 9 (2007) 90.
- [149] C. Harris et al., "Array Programming with NumPy," Nature 585 (2020) 3572, arXiv:2006.10256 [cs.MS].
- [150] P. Virtanen et al., "SciPy 1.0 Fundamental Algorithms for Scientific Computing in Python," Nat. Methods 17 (2020) 261, arXiv:1907.10121 [cs.MS].